



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

**NOVEL TUMOR ANTIGEN USEFUL IN DIAGNOSIS  
AND THERAPY OF PROSTATE AND COLON CANCER**

5

**BACKGROUND OF THE INVENTION**

Prostate cancer is the most frequently diagnosed cancer and second leading cause of cancer death in men. Some 45,000 men die annually of this disease. Only lung cancer has a higher mortality. The chance of a man developing invasive prostate cancer during his lifetime is 1 in 6. At the age of 50, a man has a greater than 40% chance of developing prostate cancer and nearly a 3% chance of dying from this disease. While some advances in the treatment of locally confined tumors have been achieved, prostate cancer is incurable once it has metastasized. Patients with metastatic prostate cancer are treated by hormonal ablation therapy, but with only short-term success. Eventually, these patients develop an androgen-refractory state leading to disease progression and death.

A continuing and fundamental problem in the management of prostate cancer is the absence of reliable diagnostic and prognostic markers capable of accurately detecting early-stage localized tumors and/or predicting disease susceptibility and progression. Early detection and diagnosis of prostate cancer currently relies on digital rectal examination (DRE), prostate specific antigen (PSA) measurements, transrectal ultrasonography (TRUS), and transrectal needle biopsy (TRNB). Serum PSA measurements in combination with DRE represent the leading diagnostic approach at present. However, this approach has major limitations which have fueled intensive research into finding better diagnostic markers of this disease. A number of markers have been identified, and at least one, PSA, is in widespread clinical use. However, ideal prostate tumor markers have been extremely elusive and no marker has yet proven reliable for predicting progression of the disease. Thus, there is a need for more reliable and informative diagnostic and prognostic methods in the management of prostate cancer.

In addition, there is also great interest in identifying prostate-specific proteins that could be appropriate as therapeutic targets, as there is no effective treatment for patients who develop recurrent disease or who have been diagnosed with metastatic disease. Although hormone ablation therapy can palliate these patients, the majority inevitably progress to develop incurable, androgen-independent disease (Lalani et al., 1997, Cancer Metastasis Rev. 16: 29-66).

PSA is the most widely used tumor marker for screening, diagnosis, and monitoring prostate cancer today. In particular, several immunoassays for the detection of serum PSA are in widespread clinical use. Recently, a reverse transcriptase-polymerase chain reaction (RT-PCR) assay for PSA mRNA in serum has been developed. However, PSA is not a disease-specific marker, as elevated levels of PSA are detectable in a large percentage of patients with BPH and prostatitis (25-86%)(Gao et al., 1997, Prostate 31: 264-281), as well as in other nonmalignant disorders and in some normal men, a factor which significantly limits the diagnostic specificity of this marker. For example, elevations in serum PSA of between 4 to 10 ng/ml are observed in BPH, and even higher values are observed in prostatitis, particularly acute prostatitis. BPH is an extremely common condition in men. Further confusing the situation is the fact that serum PSA elevations may be observed without any indication of disease from DRE, and vice-versa. Moreover, it is now recognized that PSA is not prostate-specific (Gao et al., supra, for review).

Various methods designed to improve the specificity of PSA-based detection have been described, such as measuring PSA density and the ratio of free vs. complexed PSA. However, none of these methodologies have been able to reproducibly distinguish benign from malignant prostate disease. In addition, PSA diagnostics have sensitivities of between 57-79% (Cupp & Osterling, 1993, Mayo Clin Proc 68:297-306), and thus miss identifying prostate cancer in a significant population of men with the disease.

Prostate-Specific Membrane Antigen (PSMA) is a recently described cell surface marker of prostate cancer which has been the subject of various studies evaluating its use as a diagnostic and therapeutic marker. PSMA expression is largely restricted to prostate tissues, but detectable levels of PSMA mRNA have been observed in brain, salivary gland, small intestine, and renal cell carcinoma (Israeli et al., 1993, Cancer Res 53: 227-230). PSMA protein is highly expressed in most primary and metastatic prostate cancers, but is also expressed in most intraepithelial neoplasia specimens (Gao et al., supra). Preliminary results using an Indium-111 labeled, anti-PSMA monoclonal antibody to image recurrent prostate cancer show some promise (Sodee et al., 1996, Clin Nuc Med 21: 759-766). PSMA is a hormone dependent antigen requiring the presence of functional androgen receptor. Since not all prostate cancer cells express androgen receptor, the clinical utility of PSMA as a therapeutic target may be inherently limited. Clinical trials designed to examine the effectiveness of PSMA immunotherapy are also underway.

Prostate Stem Cell Antigen (PSCA) is another very recently described cell surface marker of prostate cancer (Reiter et al., 1998, Proc. Natl. Acad. Sci. USA 95: 1735-1740). PSCA expression has been shown to be prostate specific and widely over-expressed across all stages of prostate cancer, including high grade prostatic intraepithelial neoplasia (PIN), androgen-dependent and androgen-independent prostate tumors. The PSCA gene has been mapped to chromosome 8q24.2, a region of allelic gain in more than 80% of prostate cancers. PSCA shows promise as a diagnostic and therapeutic target in view of its cell surface location, prostate specificity, and greatly upregulated expression in prostate cancer cells.

Progress in the identification of specific markers has been slow due to a lack of experimental animal model systems that recapitulate clinical disease. Attempted solutions to this problem have included the generation of prostate cancer cell lines (Horoszewicz et al., 1983, Cancer Res. 43, 1809) and prostate cancer xenografts (Pretlow et al., 1991, Cancer Res. 51, 3814; van Weerden et al., 1996, Am. J. Pathol. 149, 1055; Klein et al., 1997, Nature Med. 3, 402). However, these approaches have met with limited success. For example, xenografts have generally produced low long-term survival rates. In addition, none of the most widely used human prostate cancer cell lines - PC-3, DU-145, and LNCaP - have been shown to reproducibly give rise to osteoblastic lesions typical of prostate cancer. A further limitation of the DU-145 and PC-3 cell lines is that these cells do not express prostate specific antigen (PSA) or androgen receptor (AR) (Kaighn et al., 1979, Invest. Urol. 17: 16-23; Gleave et al., 1992, Cancer Res. 52: 1598-1605), questioning their relevance to clinical prostate cancer. The LNCaP cell line is androgen responsive and expresses PSA, but contains a mutation in the androgen receptor which alters ligand specificity.

Recently, however, a series of prostate cancer xenografts (derived from patient tumors) demonstrating genetic and phenotypic characteristics closely paralleling the human clinical situation have been described (Klein et al., 1997, Nature Med. 3: 402). These LAPC (Los Angeles Prostate Cancer) xenografts have survived passage in severe combined immune deficient (SCID) mice for longer than one year. The LAPC-4 xenograft model system has the capacity to mimic the transition from androgen dependence to androgen independence (Klein et al., 1997, supra). LAPC-4 tumors regress in male mice after castration, but re-grow within 2-3 months as androgen independent tumors. Both androgen dependent (AD) and androgen independent (AI) LAPC-4 xenograft tumors express equal levels of the prostate specific markers PSA, PSMA (prostate specific membrane antigen) and PSCA (prostate stem cell antigen), which was identified using representational difference analysis of cDNAs derived from the AD and AI variants of the LAPC-4 xenograft.

### SUMMARY OF THE INVENTION

The present invention relates to methods and compositions for the diagnosis and therapy of prostate and colon cancer, derived from or based on a novel prostate-specific, androgen-regulated, cell surface serine protease termed 20P1F12/TMPRSS2 and extensively described herein. A full length cDNA comprising the entire coding sequence of the 20P1F12/TMPRSS2 gene (also designated 20P1F12-GTC1 herein) is provided (FIG. 1). This cDNA encodes a protein which is highly related to, but structurally distinct from, the recently published TMPRSS2 (Paoloni-Giacobino et al., 1997, Genomics 44: 309-320). The 20P1F12/TMPRSS2 gene also shows a very different expression pattern relative to the expression profile of TMPRSS2.

More specifically, the invention provides polynucleotides corresponding or complementary to all or part of the 20P1F12/TMPRSS2 gene, mRNA, and/or coding sequence, preferably in isolated form, including polynucleotides encoding 20P1F12/TMPRSS2 proteins and fragments thereof, DNA, RNA, DNA/RNA hybrid, and related molecules, polynucleotides or oligonucleotides complementary to the 20P1F12/TMPRSS2 genes or mRNA sequences or parts thereof, and polynucleotides or oligonucleotides which hybridize to the 20P1F12/TMPRSS2 genes, mRNAs, or to 20P1F12/TMPRSS2-encoding polynucleotides. Also provided are means for isolating cDNAs and the genes encoding 20P1F12/TMPRSS2. Recombinant DNA molecules containing 20P1F12/TMPRSS2 polynucleotides, cells transformed or transduced with such molecules, and host-vector systems for the expression of 20P1F12/TMPRSS2 gene products are also provided. The invention further provides 20P1F12/TMPRSS2 proteins and polypeptide fragments thereof.

Methods for detecting the presence of 20P1F12/TMPRSS2 polynucleotides and proteins in various biological samples, as well as methods for identifying cells that express 20P1F12/TMPRSS2 are provided. Diagnostic imaging methods for the management of prostate and colon cancers are also provided. The invention further provides various therapeutic compositions and strategies for treating prostate cancer, including particularly, antibody therapy methods and compositions, cancer vaccines, and small molecule therapy.

The invention provides antibodies that bind to 20P1F12/TMPRSS2 proteins and polypeptide fragments thereof, including polyclonal and monoclonal antibodies, murine and other mammalian antibodies, chimeric antibodies, humanized and fully human antibodies, and antibodies labeled with a detectable marker or toxin or therapeutic composition. Several monoclonal antibodies specifically reactive with

20P1F12/TMPRSS2 are also described herein. These and other 20P1F12/TMPRSS2 antibodies are useful in molecular diagnostic assays and diagnostic imaging methods for detecting, localizing and characterizing carcinomas of the prostate and colon and metastases thereof. Cancer vaccines are also provided.

5

#### BRIEF DESCRIPTION OF THE FIGURES

FIG. 1. Nucleotide and deduced amino acid sequences of cDNA clone 20P1F12-GTC1 (Example 3)(as deposited with the ATCC; Accession No. 207097).

10

FIG. 2. Nucleotide and deduced amino acid sequences of TMPRSS2 gene sequence as published in Paoloni-Giacobino et al., 1997, Genomics 44: 309-320.

15

FIG. 3. Amino acid sequence alignment comparing 20P1F12-GTC1 cDNA (Example 3) with the previously published sequence of TMPRSS2 (Paoloni-Giacobino et al., 1997, Genomics 44: 309-320). Amino acid differences are shown in bold type.

FIG. 4. Nucleotide sequence of initially isolated SSH clone 20P1F12.

20

FIG. 5. RT-PCR analysis of 20P1F12/TMPRSS2 gene expression in prostate cancer xenografts, normal prostate, and other tissues and cell lines, showing approximately equal levels of expression in normal prostate and three prostate cancer xenografts (Panel A); and showing largely prostate specific expression in normal human tissues, with significantly lower expression levels detectable in colon, pancreas, kidney and lung (Panels B and C).

25

FIG. 6. Northern blot analysis of 20P1F12/TMPRSS2 gene expression in normal human tissues and prostate cancer xenografts using labeled clone 20P1F12 cDNA probe. Panels A and B: Expression in 16 normal tissues largely restricted to prostate; with kidney, pancreas and lung showing 10- to 20- fold lower expression levels. Panel C: Expression in LAPC-4 prostate cancer xenografts and various cell lines showing high level expression in prostate cancer xenografts, some of the prostate cancer cell lines, and in a colon carcinoma cell line. Except for LAPC-9 AI, expression of 20P1F12/TMPRSS2 in the xenografts was comparable to levels observed in normal prostate samples. In addition, lower level expression in the epidermoid carcinoma line A431 was observed.

30

35

FIG. 7. Expression of 20P1F12/TMPRSS2 in prostate and colon cancer cell lines. Xenograft and cell line filters were prepared with 10 µg of total RNA per lane. The blots

were analyzed using a 20P1F12/TMPRSS2 derived gene fragment probe. All RNA samples were normalized by ethidium bromide staining. Kilobases= kb.

FIG. 8. Characterization of Monoclonal antibodies directed against 20P1F12/TMPRSS2. Monoclonal antibodies towards 20P1F12/TMPRSS2 were generated using a purified GST-20P1F12/TMPRSS2 fusion protein as described in Example 5. Hybridoma supernatants were initially screened by ELISA against purified 20P1F12/TMPRSS2 protein cleaved from the GST-fusion. A secondary screen involved western blotting against lysates derived from 293T cells transfected with a retroviral vector encoding 20P1F12/TMPRSS2. (A) Six mAbs (1F9, 2D10, 2F8, 6B11, 8C6 and 9G8) that specifically recognize 20P1F12/TMPRSS2 were used to probe western blots from cell lysates derived from 293T cells transfected with either 20P1F12/TMPRSS2 (lane 1) or neo (as a control, lane 2). (B) Cell lysates from 293T cells transfected with 20P1F12/TMPRSS2 (lane 1) or neo (as a control, lane 2), LAPC-9 AD and LNCaP were probed with 1F9 anti-TMPRSS2 mAb. Molecular weight standards are indicated on the side in kilodaltons (KD).

FIG. 9. Cell surface biotinylation of 20P1F12/TMPRSS2. (A) His-tagged 20P1F12/TMPRSS2 or neo (as a control) were transfected into 293T cells. Intact cells were incubated with biotin to biotinylate all cell surface proteins. Cell lysates were either analyzed by western blotting directly (lanes 1 and 2, or they were incubated with streptavidin to affinity purify all labeled cell surface proteins). Streptavidin purified cell surface proteins were analyzed by western blotting using anti-His antibodies (lanes 3 and 4). Biotinylated protein was only detected in 20P1F12/TMPRSS2 transfected cells. (B) Biotinylated PC-3 (lane 2) and LNCaP (Lane 4), and unlabelled PC-3 (lane 1) and LNCaP (lane 3) were incubated with streptavidin gel and then analyzed by western blotting using 1F9 mAb. 20P1F12/TMPRSS2 was only detected in biotinylated samples. Molecular weight standards are indicated on the side in kilodaltons (KD).

FIG. 10. De-glycosylation of 20P1F12/TMPRSS2 in transfected 293T cells. His-tagged 20P1F12/TMPRSS2 transfected into 293T cells was purified using Nickel-agarose. 20P1F12/TMPRSS2 protein was then de-glycosylated using N-glycosidase F. Untreated 20P1F12/TMPRSS2 (lane 1) and de-glycosylated protein (lane 2) were analyzed by western blotting using anti-His antibodies. A shift in molecular weight is detected with de-glycosylation. Molecular weight standards are indicated on the side in kilodaltons (KD).



FIG. 11. Androgen regulation of 20P1F12/TMPRSS2 cell surface protease. LNCaP cells were deprived of androgen by growing cells in 2% charcoal-stripped fetal bovine serum for 1 week (lane 1), or 24 hours (lane 3). Androgen regulation was determined by stimulating 24 hour starved cells with 10 nM mibolerone (androgen analogue) for 9 hours (lane 4). Expression of 20P1F12/TMPRSS2 was compared to 20P1F12/TMPRSS2 levels in LNCaP cells growing in complete medium (lane 2) by northern blotting of 10  $\mu$ g of RNA/lane probed with a 20P1F12/TMPRSS2 probe. Equal RNA loading was determined by ethidium bromide staining and subsequent probing with a  $\beta$ -actin probe. PSA levels were determined as a control for androgen regulation. Molecular weight standards are indicated on the side in kilobases (kb).

FIG. 12. Androgen regulation of 20P1F12/TMPRSS2 in LNCaP. LNCaP cells were deprived of androgen by growing cells in 2% charcoal-stripped fetal bovine serum for 1 week. Androgen regulation was determined by stimulating cells with mibolerone (Mib) for various time points. Expression of 20P1F12/TMPRSS2 was determined by western blotting of cell lysates using anti-1F9 mAb. As additional controls cell lysates from PC-3 cells infected with either neo (as a control) or 20P1F12/TMPRSS2 were used. Equal protein loading was determined by probing the western blot with anti-Grb-2 antibodies (Transduction Laboratories)(data not shown). Protein expression of 20P1F12/TMPRSS2 was compared to RNA levels by northern blotting of 10  $\mu$ g RNA/lane probed with a 20P1F12/TMPRSS2 probe. Equal RNA loading was determined by probing the northern blot with a  $\beta$ -actin probe.

FIG. 13. Effect of 20P1F12/TMPRSS2 expression in NIH 3T3 cells. NIH 3T3 cells were infected with retrovirus encoding either neo (as a control) or 20P1F12/TMPRSS2. Forty-eight hours after infection the cells were analyzed by light microscopy. Cells that appeared to accumulate high numbers of vacuoles are indicated with arrows.

### DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to methods and compositions for the diagnosis and therapy of prostate cancer which utilize isolated polynucleotides corresponding to the 20P1F12/TMPRSS2 gene, proteins encoded by the 20P1F12/TMPRSS2 gene and fragments thereof, and antibodies capable of specifically recognizing and binding to 20P1F12/TMPRSS2 proteins. The 20P1F12/TMPRSS2 gene encodes a predicted 492 amino acid multimeric protein containing a serine protease domain, a scavenger receptor cysteine-rich domain, an LDL receptor class A domain, and a predicted transmembrane domain as has been described for TMPRSS2 (Paoloni-Giacobino et

al., 1997, Genomics 44: 309-320). Paoloni-Giacobino et al. found that the TMPRSS2 gene is expressed strongly in small intestine and only weakly in several other tissues and have also mapped the TMPRSS2 gene to chromosome 21. The physiological role of TMPRSS2 is unknown. Applicants have cloned a full length cDNA comprising the entire coding region of the 20P1F12/TMPRSS2 gene, but it contains several nucleotide sequence differences relative the published sequence of TMPRSS2. Five of these sequence differences result in amino acid differences. The nature and significance of these changes are presently unknown. In addition, applicants novel 20P1F12/TMPRSS2 has a completely different expression pattern in comparison to what has been known for the previously reported TMPRSS2.

Because 20P1F12/TMPRSS2 is a prostate specific protease, it is possible that it functions directly in the development and/or progression of prostate cancer, particularly in the development of metastatic disease. In this regard, proteases are known to be involved in invasion and metastasis of cancer cells (Henriet et al., 1999, APMIS 107(1):111-9; Rochefort et al., 1999, APMIS 107(1):86-95; Webber et al., 1995, Clin Cancer Res 1(10):1089-94; Duffy, 1996, Clin Cancer Res 2(4):613-8; Webber and Waghray, 1995 Clin Cancer Res 1(7):755-61). For instance, urokinase-type plasminogen activator (u-PA), cathepsin D and PSA are thought to contribute to the ability of prostate cancer cells to metastasize. The potential direct involvement of 20P1F12/TMPRSS2 function in prostate cancer, and particularly in metastasis, may be evaluated as described in Example 5.

Interestingly, the 20P1F12/TMPRSS2 and TMPRSS2 primary structure contains protein-protein interaction domains and an extracellular protease domain. The function of 20P1F12/TMPRSS2 and TMPRSS2 is unclear. The function of 20P1F12/TMPRSS2 and TMPRSS2 may involve binding to substrate proteins in the extracellular milieu through its SRCR and/ or LDLA domains. Examples of proteins that exhibit SRCR domains include: CD6, an adhesion molecule that binds to ALCAM (activated leukocyte cell adhesion molecule) and mediates thymocyte-thymic epithelium cell binding (Whitney et al., 1995, J Biol Chem 270:18187); CD5 (Ly-1) a T-cell protein that binds CD72 on B-cells and may be involved in T-B cell communication (Luo et al., 1992, J Immunol 148:1630); BSSP-3, a brain-specific serine protease with a kringle-like structure and three scavenger receptor cysteine-rich motifs (Yamamura et al., 1997, Biochem Biophys Res Commun 239:386).

The protease domain of 20P1F12/TMPRSS2 is most homologous to the protease domain of hepsin (TMPRSS1), a transmembrane serine protease that is highly

expressed in liver and up-regulated in ovarian cancer Leyus et al., 1988, *Biochemistry* 27: 1067-74; Tanimoto et al., 1997, *Cancer Res.* 57: 2884-2887).

5 The invention is based, in part, upon the isolation of a cDNA fragment corresponding to the 20P1F12/TMPRSS2 gene by Suppression Subtraction Hybridization cloning and upon the detailed molecular and biochemical characterization studies described in the Examples. The initially isolated cDNA fragment, clone 20P1F12, showed identity in an overlapping part of the 3' untranslated sequence of the recently described full length cDNA encoding TMPRSS2. Primers designed to specifically amplify the gene  
10 corresponding to 20P1F12 were then used to characterize 20P1F12/TMPRSS2 expression in prostate cancer xenografts, normal prostate, and a variety of other normal tissues. A full length cDNA comprising the entire coding sequence of the 20P1F12/TMPRSS2 gene has been isolated and sequenced and is provided herein.

15 The nucleotide and deduced amino acid sequences of the novel 20P1F12/TMPRSS2 gene (also designated 20P1F12-GTC1 herein) are shown in FIG. 1. There are significant differences in the amino acid sequences encoded by the 20P1F12-GTC1/TMPRSS2 gene compared to the previously reported sequence of TMPRSS2 (see amino acid alignment in FIG. 3). For example, four of the amino acid differences  
20 are in the protease domain, three of which are non-conservative amino acid differences and which could affect protease function and/or specificity. Applicants' novel 20P1F12/TMPRSS2 protein has been extensively characterized, as further described in the Examples sections herein. The 20P1F12/TMPRSS2 protein is a glycosylated type II transmembrane protein with an extracellular C-terminal protease  
25 domain. The 20P1F12/TMPRSS2 gene is androgen-regulated. The 20P1F12/TMPRSS2 protein is expressed on the cell surface. Expression of the 20P1F12/TMPRSS2 gene in normal tissues is prostate-specific. Expression of 20P1F12/TMPRSS2 is also observed in prostate cancer, including high level expression in advanced and metastatic disease. In addition, 20P1F12 appears to be over-expressed in colon cancer and may  
30 also be expressed in other cancers.

In addition to the differences in structure between applicants' the 20P1F12-GTC1/TMPRSS2 gene (FIG. 1) and the previously reported sequence (FIG. 2), the results of applicants expression analysis are contrary to those reported by Paoloni-  
35 Giacobino et al. In particular, applicants analysis of 20P1F12/TMPRSS2 gene expression by RT-PCR in 16 normal tissues shows the highest level expression in prostate, with substantially lower levels detected in colon, pancreas, kidney, liver and lung and no detectable expression in small intestine (FIG. 5, Panels B and C). Similar results were obtained on Northern blot analysis, although the expression level

detected in prostate by Northern blot is extremely high relative to these other tissues in which only very low level expression is detected (FIG. 6, Panels A and B).

5 Expression analysis also shows high level expression of 20P1F12/TMPRSS2 in all prostate cancer xenografts tested, at approximately the same levels seen in normal prostate (FIG. 5, Panel A). Northern blot analysis shows similar results, with somewhat lower level expression detected in the LAPC-9 xenograft relative to the LAPC-4 xenografts and normal prostate; expression is also detected in some of the prostate cancer cell lines analyzed (FIG. 6, Panel C). The 20P1F12/TMPRSS2 gene is  
10 also expressed in a number of prostate cancer cell lines (FIG. 7). These results indicate that the 20P1F12/TMPRSS2 gene is primarily a prostate specific gene which may be involved in the development and/or progression of prostate cancer. In addition, high level expression of 20P1F12/TMPRSS2 was detected by Northern blot in a number of colon carcinoma cell lines (FIG. 6, Panel C; FIG. 7). Expression of  
15 20P1F12/TMPRSS2 in colon cancer may provide a molecular basis for detecting, diagnosing, prognosing and/or treating colon cancer.

Thus, the invention provides a unique and useful 20P1F12/TMPRSS2 gene (and protein), having the nucleotide and encoded amino acid sequences as shown in FIG. 1.  
20 Nucleotide probes corresponding to all or part of the 20P1F12/TMPRSS2 cDNA sequences disclosed herein (FIGS. 1 and 4) are provided and may be used to isolate or identify other cDNAs encoding all or part of the 20P1F12/TMPRSS2 gene sequence. The invention further provided primers capable of specifically amplifying the 20P1F12/TMPRSS2 gene or its RNA transcripts. The invention further provides isolated  
25 polynucleotides containing coding sequences of the 20P1F12/TMPRSS2 gene product(s). Such polynucleotides may be used to express 20P1F12/TMPRSS2 encoded proteins and peptides having a number of further uses. 20P1F12/TMPRSS2 gene probes and primers may also be used to detect the presence or absence of 20P1F12/TMPRSS2 mRNA in various biological samples, for detecting prostate cancer cells and other cells expressing 20P1F12/TMPRSS2, for generating tumor vaccines, and in molecular diagnostic and prognostic assays for prostate cancer. Polynucleotides corresponding or complementary to the 20P1F12/TMPRSS2 gene may be useful in methods for treating prostate cancer, such as, for example, in modulating or inhibiting  
30 20P1F12/TMPRSS2 biological activity.

35 More specifically, a 20P1F12/TMPRSS2 polynucleotide useful in the practice of the invention may comprise a polynucleotide having the nucleotide sequence of human 20P1F12/TMPRSS2 as shown in FIG. 1 (SEQ ID NO. XX) or the nucleotide sequence of the previously reported TMPRSS2 as shown in FIG. 2 (SEQ ID NO: XX), a sequence

complementary to either of the foregoing, or a polynucleotide fragment of any of the foregoing. Another embodiment comprises a polynucleotide which encodes the 20P1F12/TMPRSS2 protein amino acid sequence as shown in FIG. 1 (SEQ ID NO. XX), a sequence complementary thereto, or a polynucleotide fragment thereof. Another  
5 embodiment comprises a polynucleotide which is capable of hybridizing under stringent hybridization conditions to the 20P1F12/TMPRSS2 cDNA shown in FIG. 1 (SEQ ID NO. XX) or to a polynucleotide fragment thereof.

10 Included within the scope of this aspect of the invention are genomic DNA, cDNAs, ribozymes, and antisense molecules, as well as nucleic acid molecules based on an alternative backbone or including alternative bases, whether derived from natural sources or synthesized. For example, antisense molecules can be RNAs or other  
15 molecules, including peptide nucleic acids (PNAs) or non-nucleic acid molecules such as phosphorothioate derivatives, that specifically bind DNA or RNA in a base pair-dependent manner. A skilled artisan can readily obtain these classes of nucleic acid molecules using the 20P1F12/TMPRSS2 polynucleotides and polynucleotide sequences disclosed herein.

20 Further specific embodiments of this aspect of the invention include primers and primer pairs, which allow the specific amplification of the polynucleotides of the invention or of any specific parts thereof, and probes that selectively or specifically hybridize to nucleic acid molecules of the invention or to any part thereof. Probes may be labeled with a detectable marker, such as, for example, a radioisotope, fluorescent  
25 compound, bioluminescent compound, a chemiluminescent compound, metal chelator or enzyme. Such probes and primers can be used to detect the presence of a 20P1F12/TMPRSS2 polynucleotide in a sample and as a means for detecting a cell expressing a 20P1F12/TMPRSS2 protein. An example of such a probe is a polypeptide comprising all or part of the human 20P1F12/TMPRSS2 cDNA sequence shown in FIG. 1 (SEQ ID NO. XX). Examples of primer pairs capable of specifically amplifying  
30 20P1F12/TMPRSS2 mRNAs are also described in the Examples which follow. As will be understood by the skilled artisan, a great many different primers and probes may be prepared based on the sequences provided in herein and used effectively to amplify and/or detect a 20P1F12/TMPRSS2 mRNA.

35 The 20P1F12/TMPRSS2 polynucleotides of the invention are useful for a variety of purposes, including but not limited to their use as probes and primers for the amplification and/or detection of the 20P1F12/TMPRSS2 gene, mRNA, or fragments thereof; as reagents for the diagnosis and/or prognosis of prostate and colon cancer; as coding sequences capable of directing the expression of 20P1F12/TMPRSS2

polypeptides; as tools for modulating or inhibiting the expression of the 20P1F12/TMPRSS2 gene and/or translation of the 20P1F12/TMPRSS2 transcript; and as therapeutic agents.

- 5 The invention also provides 20P1F12/TMPRSS2 proteins and polypeptides which may be used, for example, to generate antibodies or for use as cancer vaccines. Antibodies capable of specifically binding to and identifying 20P1F12/TMPRSS2 proteins or polypeptides may be used to detect the expression of 20P1F12/TMPRSS2, determine its subcellular location, detect and image prostate cancer cells and prostate tumors, and  
10 modulate or inhibit 20P1F12/TMPRSS2 biological activity. Antibodies may also be used therapeutically as described further below. Methods for the generation of polyclonal and monoclonal antibodies are well known in the art.

- The invention also provides recombinant DNA or RNA molecules containing a  
15 20P1F12/TMPRSS2 polynucleotide, including but not limited to phages, plasmids, phagemids, cosmids, YACs, BACs, as well as various viral and non-viral vectors well known in the art, and cells transformed or transfected with such recombinant DNA or RNA molecules. As used herein, a recombinant DNA or RNA molecule is a DNA or RNA molecule that has been subjected to molecular manipulation in vitro. Methods for  
20 generating such molecules are well known (see, for example, Sambrook et al, 1989, supra).

- The invention further provides a host-vector system comprising a recombinant DNA molecule containing a 20P1F12/TMPRSS2 polynucleotide within a suitable prokaryotic  
25 or eukaryotic host cell. Examples of suitable eukaryotic host cells include a yeast cell, a plant cell, or an animal cell, such as a mammalian cell or an insect cell (e.g., a baculovirus-infectible cell such as an Sf9 cell). Examples of suitable mammalian cells include various prostate cancer cell lines such as LnCaP, PC-3, DU145, LAPC-4, TsuPr1, other transfectable or transducible prostate cancer cell lines, as well as a number of  
30 mammalian cells routinely used for the expression of recombinant proteins (e.g., COS, CHO, 293, 293T cells). More particularly, a polynucleotide comprising the coding sequence of a 20P1F12/TMPRSS2 may be used to generate 20P1F12/TMPRSS2 proteins or fragments thereof using any number of host-vector systems routinely used and widely known in the art.

- 35 A wide range of host-vector systems suitable for the expression of 20P1F12/TMPRSS2 proteins or fragments thereof are available, see for example, Sambrook et al., 1989, supra; Current Protocols in Molecular Biology, 1995, supra). Preferred vectors for mammalian expression include but are not limited to pcDNA 3.1 myc-His-tag

(Invitrogen) and the retroviral vector pSR $\alpha$ kneo (Muller et al., 1991, MCB 11:1785). Using these expression vectors, 20P1F12/TMPRSS2 may be preferably expressed in several prostate and non-prostate cancer cell lines, including for example 3T3, 293, 293TPC-3, LNCaP and TsuPr1. The host-vector systems of the invention are useful for the production of a 20P1F12/TMPRSS2 protein or fragment thereof. Such host-vector systems may be employed to study the functional properties of 20P1F12/TMPRSS2 and 20P1F12/TMPRSS2 mutations.

Proteins encoded by the 20P1F12/TMPRSS2 genes, or by fragments thereof, will have a variety of uses, including but not limited to generating antibodies and in methods for identifying ligands and other agents and cellular constituents that bind to a 20P1F12/TMPRSS2 gene product. Such proteins may also be used as cancer vaccines. Antibodies raised against a 20P1F12/TMPRSS2 protein or fragment thereof may be useful in diagnostic and prognostic assays, imaging methodologies (including, particularly, cancer imaging), and therapeutic methods in the management of human cancers characterized by expression of a 20P1F12/TMPRSS2 protein, such as prostate and colon cancers. Various immunological assays useful for the detection of 20P1F12/TMPRSS2 proteins are contemplated, including but not limited to various types of radioimmunoassays, enzyme-linked immunosorbent assays (ELISA), enzyme-linked immunofluorescent assays (ELIFA), immunocytochemical methods, and the like. Such antibodies may be labeled and used as immunological imaging reagents capable of detecting prostate cells (e.g., in radioscintigraphic imaging methods).

In a specific embodiment, a novel 20P1F12/TMPRSS2 protein having the amino acid sequence of human 20P1F12/TMPRSS2 is provided in FIG. 1 (SEQ ID NO. XX). Fusion proteins which combine all or part of 20P1F12/TMPRSS2 with a heterologous polypeptide are also contemplated. The 20P1F12/TMPRSS2 protein of the invention may be embodied in many forms, preferably in isolated form. As used herein, the protein is said to be "isolated" when physical, mechanical or chemical methods are employed to remove the 20P1F12/TMPRSS2 protein from cellular constituents that are normally associated with the protein. A skilled artisan can readily employ standard purification methods to obtain an isolated 20P1F12/TMPRSS2 protein. A purified 20P1F12/TMPRSS2 protein molecule will be substantially free of other proteins or molecules which impair the binding of 20P1F12/TMPRSS2 to antibody or other ligand. The nature and degree of isolation and purification will depend on the intended use. Embodiments of a 20P1F12/TMPRSS2 protein include a purified 20P1F12/TMPRSS2 protein and a functional, soluble 20P1F12/TMPRSS2 protein. In one form, such functional, soluble 20P1F12/TMPRSS2 proteins or fragments thereof retain the ability to bind antibody or other ligand.

- Recombinant methods can be used to generate nucleic acid molecules that encode the 20P1F12/TMPRSS2 protein. In this regard, the 20P1F12/TMPRSS2-encoding nucleic acid molecules described herein provide means for generating defined fragments of the 20P1F12/TMPRSS2 protein. Such 20P1F12/TMPRSS2 polypeptides are particularly useful in generating domain specific antibodies (e.g., antibodies recognizing an extracellular epitope of the 20P1F12/TMPRSS2 protein), identifying agents or cellular factors that bind to a particular 20P1F12/TMPRSS2 domain, and in various therapeutic contexts, including but not limited to cancer vaccines. 20P1F12/TMPRSS2 polypeptides containing particularly interesting structures can be predicted and/or identified using various analytical techniques well known in the art, including, for example, the methods of Chou-Fasman, Garnier-Robson, Kyte-Doolittle, Eisenberg, Karplus-Schultz or Jameson-Wolf analysis, or on the basis of immunogenicity.
- Another aspect of the invention provides antibodies that immunospecifically bind to the 20P1F12/TMPRSS2 protein and polypeptide fragments thereof. The most preferred antibodies will selectively bind to a 20P1F12/TMPRSS2 protein and will not bind (or will bind weakly) to non-20P1F12/TMPRSS2 proteins and polypeptides. Anti-20P1F12/TMPRSS2 antibodies that are particularly contemplated include monoclonal and polyclonal antibodies as well as fragments containing the antigen binding domain and/or one or more complementarity determining regions of these antibodies. As used herein, an antibody fragment is defined as at least a portion of the variable region of the immunoglobulin molecule which binds to its target, i.e., the antigen binding region.
- For some applications, it may be desirable to generate antibodies which specifically react with a particular 20P1F12/TMPRSS2 protein and/or an epitope within a particular structural domain. For example, preferred antibodies useful for cancer diagnostic imaging purposes are those which react with an epitope in an extracellular region of the 20P1F12/TMPRSS2 protein as expressed in cancer cells. Such antibodies may be generated by using the 20P1F12/TMPRSS2 protein, or using peptides derived from predicted extracellular or other domains of 20P1F12/TMPRSS2, and used as an immunogen.
- The 20P1F12/TMPRSS2 antibodies of the invention may be particularly useful in prostate and colon cancer diagnostic and prognostic assays, imaging methodologies, and therapeutic strategies. The invention provides various immunological assays useful for the detection and quantification of 20P1F12/TMPRSS2. Such assays generally comprise one or more 20P1F12/TMPRSS2 antibodies capable of recognizing and binding a 20P1F12/TMPRSS2, and may be performed within various immunological assay



formats well known in the art, including but not limited to various types of radioimmunoassays, enzyme-linked immunosorbent assays (ELISA), enzyme-linked immunofluorescent assays (ELIFA), and the like. In addition, immunological imaging methods capable of detecting prostate cancer are also provided by the invention, including but limited to radioscintigraphic imaging methods using labeled 20P1F12/TMPRSS2 antibodies. Such assays may be clinically useful in the detection, monitoring, and prognosis of prostate cancer, particularly advanced prostate cancer.

20P1F12/TMPRSS2 antibodies may also be used in methods for purifying 20P1F12/TMPRSS2 proteins and polypeptides and for isolating 20P1F12/TMPRSS2 homologues and related molecules. For example, in one embodiment, the method of purifying a 20P1F12/TMPRSS2 protein comprises incubating a 20P1F12/TMPRSS2 antibody, which has been coupled to a solid matrix, with a lysate or other solution containing 20P1F12/TMPRSS2 under conditions which permit the 20P1F12/TMPRSS2 antibody to bind to 20P1F12/TMPRSS2; washing the solid matrix to eliminate impurities; and eluting the 20P1F12/TMPRSS2 from the coupled antibody. Other uses of the 20P1F12/TMPRSS2 antibodies of the invention include generating anti-idiotypic antibodies that mimic the 20P1F12/TMPRSS2 protein.

20P1F12/TMPRSS2 antibodies may also be used therapeutically by, for example, modulating or inhibiting the biological activity of a 20P1F12/TMPRSS2 protein or targeting and destroying prostate cancer cells expressing a 20P1F12/TMPRSS2 protein. Antibody therapy of prostate and colon cancer is described in further detail below.

Various methods for the preparation of antibodies are well known in the art. For example, antibodies may be prepared by immunizing a suitable mammalian host using a 20P1F12/TMPRSS2 protein, peptide, or fragment, in isolated or immunoconjugated form (Antibodies: A Laboratory Manual, CSH Press, Eds., Harlow, and Lane (1988); Harlow, Antibodies, Cold Spring Harbor Press, NY (1989)). In addition, fusion proteins of 20P1F12/TMPRSS2 may also be used, such as a 20P1F12/TMPRSS2 GST-fusion protein. In a particular embodiment, a GST fusion protein comprising all or most of the open reading frame amino acid sequence of FIG. 1 may be produced and used as an immunogen to generate appropriate antibodies. As described in Example 5, such a GST fusion was used to generate several monoclonal antibodies which immunospecifically react with 20P1F12/TMPRSS2. Cells expressing or overexpressing 20P1F12/TMPRSS2 may also be used for immunizations. Similarly, any cell engineered to express 20P1F12/TMPRSS2 may be used. This strategy may result in the production of monoclonal antibodies with enhanced capacities for recognizing endogenous

20P1F12/TMPRSS2. Additional strategies for generating 20P1F12/TMPRSS2 antibodies are described in Example 5 herein.

5 The amino acid sequence of 20P1F12/TMPRSS2 as shown in FIG. 1 (SEQ ID NO. XX) may be used to select specific regions of the 20P1F12/TMPRSS2 protein for generating antibodies. For example, hydrophobicity and hydrophilicity analyses of the 20P1F12/TMPRSS2 amino acid sequence may be used to identify hydrophilic regions in the 20P1F12/TMPRSS2 structure. Regions of the 20P1F12/TMPRSS2 protein that show immunogenic structure, as well as other regions and domains, can readily be identified  
10 using various other methods known in the art, such as Chou-Fasman, Garnier-Robson, Kyte-Doolittle, Eisenberg, Karplus-Schultz or Jameson-Wolf analysis.

Methods for preparing a protein or polypeptide for use as an immunogen and for preparing immunogenic conjugates of a protein with a carrier such as BSA, KLH, or  
15 other carrier proteins are well known in the art. In some circumstances, direct conjugation using, for example, carbodiimide reagents may be used; in other instances linking reagents such as those supplied by Pierce Chemical Co., Rockford, IL, may be effective. Administration of a 20P1F12/TMPRSS2 immunogen is conducted generally by injection over a suitable time period and with use of a suitable adjuvant, as is generally  
20 understood in the art. During the immunization schedule, titers of antibodies can be taken to determine adequacy of antibody formation.

Anti-20P1F12/TMPRSS2 monoclonal antibodies are preferred and may be produced by various means well known in the art. For example, immortalized cell lines which secrete  
25 a desired monoclonal antibody may be prepared using the standard method of Kohler and Milstein or modifications which effect immortalization of lymphocytes or spleen cells, as is generally known. The immortalized cell lines secreting the desired antibodies are screened by immunoassay in which the antigen is the 20P1F12/TMPRSS2 protein or 20P1F12/TMPRSS2 fragment. When the appropriate immortalized cell culture secreting  
30 the desired antibody is identified, the cells may be expanded and antibodies produced either from in vitro cultures or from ascites fluid.

The antibodies or fragments may also be produced, using current technology, by recombinant means. Regions that bind specifically to the desired regions of the  
35 20P1F12/TMPRSS2 protein can also be produced in the context of chimeric or CDR grafted antibodies of multiple species origin. Humanized or human 20P1F12/TMPRSS2 antibodies may also be produced and are preferred. Various approaches for producing such humanized antibodies are known, and include chimeric and CDR grafting methods; methods for producing fully human monoclonal antibodies include phage display and

transgenic methods (for review, see Vaughan et al., 1998, Nature Biotechnology 16: 535-539).

5 Fully human 20P1F12/TMPRSS2 monoclonal antibodies may be generated using cloning technologies employing large human Ig gene combinatorial libraries (i.e., phage display)(Griffiths and Hoogenboom, Building an in vitro immune system: human antibodies from phage display libraries. In: Protein Engineering of Antibody Molecules for Prophylactic and Therapeutic Applications in Man. Clark, M. (Ed.), Nottingham Academic, pp 45-64 (1993); Burton and Barbas, Human Antibodies from combinatorial  
10 libraries. *Id.*, pp 65-82). Fully human 20P1F12/TMPRSS2 monoclonal antibodies may also be produced using transgenic mice engineered to contain human immunoglobulin gene loci as described in PCT Patent Application WO98/24893, Jakobovits et al., published December 3, 1997 (see also, Kucherlapati and Jakobovits, 1998, Exp. Opin. Invest. Drugs 7(4): 607-614). This method avoids the in vitro manipulation required with  
15 phage display technology and efficiently produces high affinity authentic human antibodies.

Reactivity of 20P1F12/TMPRSS2 antibodies with a 20P1F12/TMPRSS2 protein may be established by a number of well known means, including Western blot,  
20 immunoprecipitation, ELISA, and FACS analyses using, as appropriate, 20P1F12/TMPRSS2 proteins, peptides, 20P1F12/TMPRSS2-expressing cells or extracts thereof.

A 20P1F12/TMPRSS2 antibody or fragment thereof of the invention may be labeled  
25 with a detectable marker or conjugated to a second molecule, such as a cytotoxic or therapeutic agent, and used for targeting a 20P1F12/TMPRSS2 positive cell (Vitetta, E.S. et al., 1993, Immunotoxin therapy, in DeVita, Jr., V.T. et al., eds, Cancer: Principles and Practice of Oncology, 4th ed., J.B. Lippincott Co., Philadelphia, 2624-2636). Suitable detectable markers include, but are not limited to, a radioisotope, a  
30 fluorescent compound, a bioluminescent compound, chemiluminescent compound, a metal chelator or an enzyme.

The 20P1F12/TMPRSS2 protein is a cell surface serine protease that may be involved in invasion and metastasis of prostate and colon cancer. Accordingly,  
35 20P1F12/TMPRSS2 may be ideal target for therapeutic intervention. Its extracellular protease domain may be a potential drug target, while the whole extracellular domain may be a potential therapeutic antibody target. Therefore, the invention provides various immunotherapeutic compositions and methods for treating prostate and colon cancer, including antibody therapy, in vivo vaccines, and ex vivo immunotherapy

approaches, which utilize polynucleotides and polypeptides corresponding to 20P1F12/TMPRSS2 and anti-20P1F12/TMPRSS2 antibodies.

5 In one approach, anti-20P1F12/TMPRSS2 antibodies may be used to treat prostate and colon cancer. For example, unconjugated anti-20P1F12/TMPRSS2 antibody may be introduced into a patient such that the antibody binds to 20P1F12/TMPRSS2 on prostate or colon cancer cells and mediates the destruction of the cells and the tumor. The therapeutic mechanism of action may include complement-mediated cytotoxicity, antibody-dependent cellular cytotoxicity, altering the physiologic function of 20P1F12/TMPRSS2, 10 and/or the inhibition of ligand binding or signal transduction pathways. Anti-20P1F12/TMPRSS2 antibodies conjugated to toxic agents such as ricin, or to therapeutic agents, may also be used therapeutically to deliver the toxic or therapeutic agent directly to 20P1F12/TMPRSS2-bearing prostate tumor cells and thereby destroy the tumor.

15 Prostate cancer immunotherapy using anti-20P1F12/TMPRSS2 antibodies may follow the teachings generated from various approaches which have been successfully employed with respect to other types of cancer, including but not limited to colon cancer (Arlen et al., 1998, Crit Rev Immunol 18: 133-138), multiple myeloma (Ozaki et al., 1997, 20 Blood 90: 3179-3186; Tsunenari et al., 1997, Blood 90: 2437-2444), gastric cancer (Kasprzyk et al., 1992, Cancer Res 52: 2771-2776), B-cell lymphoma (Funakoshi et al., 1996, J Immunother Emphasis Tumor Immunol 19: 93-101), leukemia (Zhong et al., 1996, Leuk Res 20: 581-589), colorectal cancer (Moun et al., 1994, Cancer Res 54: 6160-6166); Velders et al., 1995, Cancer Res 55: 4398-4403), and breast cancer (Shepard et al., 25 1991, J Clin Immunol 11: 117-127).

20P1F12/TMPRSS2 antibodies may be introduced into a patient such that the antibody binds to 20P1F12/TMPRSS2 on the cancer cells and mediates the destruction of the cells and the tumor and/or inhibits the growth of the cells or the tumor. Mechanisms by which 30 such antibodies exert a therapeutic effect may include complement-mediated cytotoxicity, antibody-dependent cellular cytotoxicity, modulating the physiologic function of 20P1F12/TMPRSS2, inhibiting ligand binding or signal transduction pathways, modulating tumor cell differentiation, altering tumor angiogenesis factor profiles, and/or by inducing apoptosis. 20P1F12/TMPRSS2 antibodies conjugated to toxic or therapeutic 35 agents may also be used therapeutically to deliver the toxic or therapeutic agent directly to 20P1F12/TMPRSS2-bearing tumor cells.

Although 20P1F12/TMPRSS2 antibody therapy may be useful for all stages of the foregoing cancers, antibody therapy may be particularly appropriate in advanced or

metastatic prostate and colon cancers. In particular, because the 20P1F12/TMPRSS2 gene appears not to be regulated by androgen, anti-20P1F12/TMPRSS2 antibody therapy may be used to treat patients undergoing androgen ablation therapy. Combining the antibody therapy method of the invention with a chemotherapeutic regimen may be preferred in patients who have not received chemotherapeutic treatment, whereas treatment with the antibody therapy of the invention may be indicated for patients who have received one or more chemotherapy. Additionally, antibody therapy may also enable the use of reduced dosages of concomitant chemotherapy, particularly in patients that do not tolerate the toxicity of the chemotherapeutic agent very well.

It may be desirable for patients to be evaluated for the presence and level of 20P1F12/TMPRSS2, preferably using immunohistochemical assessments of tumor tissue, quantitative 20P1F12/TMPRSS2 imaging, or other techniques capable of reliably indicating the presence and degree of expression. Immunohistochemical analysis of tumor biopsies or surgical specimens may be preferred for this purpose. Methods for immunohistochemical analysis of tumor tissues are well known in the art.

Anti-20P1F12/TMPRSS2 monoclonal antibodies useful in treating prostate and other cancers include those which are capable of initiating a potent immune response against the tumor and those which are capable of direct cytotoxicity. In this regard, anti-20P1F12/TMPRSS2 mAbs may elicit tumor cell lysis by either complement-mediated or antibody-dependent cell cytotoxicity (ADCC) mechanisms, both of which require an intact Fc portion of the immunoglobulin molecule for interaction with effector cell Fc receptor sites or complement proteins. In addition, anti-20P1F12/TMPRSS2 mAbs which exert a direct biological effect on tumor growth are useful in the practice of the invention. Potential mechanisms by which such directly cytotoxic mAbs may act include inhibition of cell growth, modulation of cellular differentiation, modulation of tumor angiogenesis factor profiles, and the induction of apoptosis. The mechanism by which a particular anti-20P1F12/TMPRSS2 mAb exerts an anti-tumor effect may be evaluated using any number of in vitro assays designed to determine ADCC and complement-mediated cell lysis, as well as growth inhibition, modulation of apoptosis and inhibition of differentiation, and/or inhibition of angiogenesis, as is generally known in the art.

The anti-tumor activity of a particular anti-20P1F12/TMPRSS2 mAb, or combination of anti-20P1F12/TMPRSS2 mAbs, may be evaluated in vivo using a suitable animal model. For example, xenogenic prostate cancer models wherein human prostate cancer explants or passaged xenograft tissues are introduced into immune compromised animals, such as nude or SCID mice, are appropriate in relation to prostate cancer and

have been described (Klein et al., 1997, Nature Medicine 3: 402-408). For Example, PCT Patent Application WO98/16628, Sawyers et al., published April 23, 1998, describes various xenograft models of human prostate cancer capable of recapitulating the development of primary tumors, micrometastasis, and the formation of osteoblastic metastases characteristic of late stage disease. Efficacy may be predicted using inhibition of tumor formation, tumor regression, metastasis, and the like.

It should be noted that the use of murine or other non-human monoclonal antibodies, human/mouse chimeric mAbs may induce moderate to strong immune responses in some patients. In the most severe cases, such an immune response may lead to the extensive formation of immune complexes which, potentially, can cause renal failure. Accordingly, preferred monoclonal antibodies used in the practice of the therapeutic methods of the invention are those which are either fully human or humanized and which bind specifically to the target 20P1F12/TMPRSS2 antigen with high affinity but exhibit low or no antigenicity in the patient.

The method of the invention contemplate the administration of single anti-20P1F12/TMPRSS2 mAbs as well as combinations, or "cocktails, of different mAbs. Such mAb cocktails may have certain advantages inasmuch as they contain mAbs which exploit different epitope specificity, different effector mechanisms, or combine directly cytotoxic mAbs with mAbs that rely on immune effector functionality. Such mAbs in combination may exhibit synergistic therapeutic effects. In addition, the administration of anti-20P1F12/TMPRSS2 mAbs may be combined with other therapeutic agents or radiation therapy, including but not limited to various chemotherapeutic agents, androgen-blockers, and immune modulators (e.g., IL-2, GM-CSF). The anti-20P1F12/TMPRSS2 mAbs may be administered in their "naked" or unconjugated form, or may have therapeutic or toxic agents conjugated to them.

The anti-20P1F12/TMPRSS2 monoclonal antibodies used in the practice of the method of the invention may be formulated into pharmaceutical compositions comprising a carrier suitable for the desired delivery method. Suitable carriers include any material which when combined with the anti-20P1F12/TMPRSS2 mAbs retains the specificity and anti-tumor function of the antibody and is non-reactive with the subject's immune systems. Examples include, but are not limited to, any of a number of standard pharmaceutical carriers such as sterile phosphate buffered saline solutions, bacteriostatic water, and the like.

The anti-20P1F12/TMPRSS2 antibody formulations may be administered via any route capable of delivering the antibodies to the tumor site. Potentially effective routes of

administration include, but are not limited to, intravenous, intraperitoneal, intramuscular, intratumor, intradermal, and the like. The preferred route of administration is by intravenous injection. A preferred formulation for intravenous injection comprises the anti-20P1F12/TMPRSS2 mAbs in a solution of preserved bacteriostatic water, sterile unpreserved water, and/or diluted in polyvinylchloride or polyethylene bags containing 0.9% sterile Sodium Chloride for Injection, USP. The anti-20P1F12/TMPRSS2 mAb preparation may be lyophilized and stored as a sterile powder, preferably under vacuum, and then reconstituted in bacteriostatic water containing, for example, benzyl alcohol preservative, or in sterile water prior to injection.

Treatment will generally involve the repeated administration of the anti-20P1F12/TMPRSS2 antibody preparation via an acceptable route of administration such as intravenous injection (IV), typically at a dose in the range of about 0.1 to about 10 mg/kg body weight. Doses in the range of 10-500 mg mAb per week may be effective and well tolerated. Based on clinical experience with the Herceptin mAb in the treatment of metastatic breast cancer, an initial loading dose of approximately 4 mg/kg patient body weight IV followed by weekly doses of about 2 mg/kg IV of the anti-20P1F12/TMPRSS2 mAb preparation may represent an acceptable dosing regimen. Preferably, the initial loading dose is administered as a 90 minute or longer infusion. The periodic maintenance dose may be administered as a 30 minute or longer infusion, provided the initial dose was well tolerated. However, as one of skill in the art will understand, various factors will influence the ideal dose regimen in a particular case. Such factors may include, for example, the binding affinity and half life of the mAb or mAbs used, the degree of 20P1F12/TMPRSS2 overexpression in the patient, the extent of circulating shed 20P1F12/TMPRSS2 antigen, the desired steady-state antibody concentration level, frequency of treatment, and the influence of chemotherapeutic agents used in combination with the treatment method of the invention.

Optimally, patients should be evaluated for the level of circulating shed 20P1F12/TMPRSS2 antigen in serum in order to assist in the determination of the most effective dosing regimen and related factors. Such evaluations may also be used for monitoring purposes throughout therapy, and may be useful to gauge therapeutic success in combination with evaluating other parameters (such as serum PSA levels in prostate cancer therapy).

The invention further provides prostate cancer vaccines comprising a 20P1F12/TMPRSS2 protein or fragment thereof. The use of a tumor antigen in a vaccine for generating humoral and cell-mediated immunity for use in anti-cancer therapy is well

known in the art and has been employed in prostate cancer using human PSMA and rodent PAP immunogens (Hodge et al., 1995, Int. J. Cancer 63: 231-237; Fong et al., 1997, J. Immunol. 159: 3113-3117). Such methods can be readily practiced by employing a 20P1F12/TMPRSS2 protein, or fragment thereof, or a 20P1F12/TMPRSS2 -  
5 encoding nucleic acid molecule and recombinant vectors capable of expressing and appropriately presenting the 20P1F12/TMPRSS2 immunogen.

For example, viral gene delivery systems may be used to deliver a 20P1F12/TMPRSS2 - encoding nucleic acid molecule. Various viral gene delivery systems which can be used  
10 in the practice of this aspect of the invention include, but are not limited to, vaccinia, fowlpox, canarypox, adenovirus, influenza, poliovirus, adeno-associated virus, lentivirus, and sindbus virus (Restifo, 1996, Curr. Opin. Immunol. 8: 658-663). Non-viral delivery systems may also be employed by using naked DNA encoding a 20P1F12/TMPRSS2 protein or fragment thereof introduced into the patient (e.g.,  
15 intramuscularly) to induce an anti-tumor response. In one embodiment, the full-length human 20P1F12/TMPRSS2 cDNA may be employed. In another embodiment, 20P1F12/TMPRSS2 nucleic acid molecules encoding specific cytotoxic T lymphocyte (CTL) epitopes may be employed. CTL epitopes can be determined using specific algorithms (e.g., Epimer, Brown University) to identify peptides within a  
20 20P1F12/TMPRSS2 protein which are capable of optimally binding to specified HLA alleles.

Various ex vivo strategies may also be employed. One approach involves the use of dendritic cells to present 20P1F12/TMPRSS2 antigen to a patient's immune system.  
25 Dendritic cells express MHC class I and II, B7 costimulator, and IL-12, and are thus highly specialized antigen presenting cells. In prostate cancer, autologous dendritic cells pulsed with peptides of the prostate-specific membrane antigen (PSMA) are being used in a Phase I clinical trial to stimulate prostate cancer patients' immune systems (Tjoa et al., 1996, Prostate 28: 65-69; Murphy et al., 1996, Prostate 29: 371-  
30 380). Dendritic cells can be used to present 20P1F12/TMPRSS2 peptides to T cells in the context of MHC class I and II molecules. In one embodiment, autologous dendritic cells are pulsed with 20P1F12/TMPRSS2 peptides capable of binding to MHC molecules. In another embodiment, dendritic cells are pulsed with the complete 20P1F12/TMPRSS2 protein. Yet another embodiment involves engineering the  
35 overexpression of the 20P1F12/TMPRSS2 gene in dendritic cells using various implementing vectors known in the art, such as adenovirus (Arthur et al., 1997, Cancer Gene Ther. 4: 17-25), retrovirus (Henderson et al., 1996, Cancer Res. 56: 3763-3770), lentivirus, adeno-associated virus, DNA transfection (Ribas et al., 1997,



Cancer Res. 57: 2865-2869), and tumor-derived RNA transfection (Ashley et al., 1997, J. Exp. Med. 186: 1177-1182).

5 Anti-idiotypic anti-20P1F12/TMPRSS2 antibodies can also be used in anti-cancer therapy as a vaccine for inducing an immune response to cells expressing a 20P1F12/TMPRSS2 protein. Specifically, the generation of anti-idiotypic antibodies is well known in the art and can readily be adapted to generate anti-idiotypic anti-20P1F12/TMPRSS2 antibodies that mimic an epitope on a 20P1F12/TMPRSS2 protein (see, for example, Wagner et al., 1997, Hybridoma 16: 33-40; Foon et al., 1995, J Clin Invest 96: 334-342; Herlyn et al., 1996, Cancer Immunol Immunother 43: 65-76). Such an  
10 anti-idiotypic antibody can be used in anti-idiotypic therapy as presently practiced with other anti-idiotypic antibodies directed against tumor antigens.

15 Genetic immunization methods may be employed to generate prophylactic or therapeutic humoral and cellular immune responses directed against cancer cells expressing 20P1F12/TMPRSS2, particularly colon and prostate cancer cells. Constructs comprising DNA encoding a 20P1F12/TMPRSS2 protein/immunogen and appropriate regulatory sequences may be injected directly into muscle or skin of an individual, such that the cells of the muscle or skin take-up the construct and express  
20 the encoded 20P1F12/TMPRSS2 protein/immunogen. The 20P1F12/TMPRSS2 protein/immunogen may be expressed as a cell surface protein or be secreted. Expression of the 20P1F12/TMPRSS2 protein/immunogen results in the generation of prophylactic or therapeutic humoral and cellular immunity against prostate cancer. Various prophylactic and therapeutic genetic immunization techniques known in the art  
25 may be used (for review, see information and references published at internet address [www.genweb.com](http://www.genweb.com)).

Another aspect of the invention is directed to molecular diagnostic and diagnostic imaging methods which utilize the 20P1F12/TMPRSS2 polynucleotides and antibodies  
30 described herein. The expression profile and cell surface localization of 20P1F12/TMPRSS2 makes it a potential imaging reagent for metastasized disease. 20P1F12/TMPRSS2 is expressed in various prostate cancer xenograft tissues and cell lines, and is also expressed in some colon cancer cell lines. The expression status of 20P1F12/TMPRSS2 may provide information useful for localizing tumors, predicting  
35 susceptibility to advanced stage disease, and/or gauging tumor aggressiveness. 20P1F12/TMPRSS2 expression status in patient samples may be analyzed by, for example: (i) immunohistochemical analysis, (ii) in situ hybridization, (iii) RT-PCR analysis on laser capture micro-dissected samples, (iv) western blot analysis of clinical samples and cell lines, (v) tissue array analysis, (vi) in vivo imaging. Various immunological

- assays useful for the detection of 20P1F12/TMPRSS2 proteins are contemplated, including but not limited to various types of radioimmunoassays, enzyme-linked immunosorbent assays (ELISA), enzyme-linked immunofluorescent assays (ELIFA), immunocytochemical methods, and the like. As an example, 20P1F12/TMPRSS2 antibodies may be labeled and used as immunological imaging reagents capable of detecting prostate and colon cancer cells (e.g., in radioscintigraphic imaging methods). For radioscintigraphic in vivo imaging, radiolabeled 20P1F12/TMPRSS2 antibodies specifically reactive with extracellular epitopes of 20P1F12/TMPRSS2 are preferred.
- 10 Assays for identifying prostate, prostate cancer or colon cancer cells comprise detecting polynucleotides corresponding to the 20P1F12/TMPRSS2 gene in a biological sample, such as serum, bone, prostate, colon and other tissues, urine, semen, cell preparations, and the like. Detectable 20P1F12/TMPRSS2 polynucleotides include, for example, a 20P1F12/TMPRSS2 gene or fragments thereof, 20P1F12/TMPRSS2 mRNA, 15 alternative splice variant 20P1F12/TMPRSS2 mRNAs, and recombinant DNA or RNA molecules containing a 20P1F12/TMPRSS2 polynucleotide. A number of methods for amplifying and/or detecting the presence of 20P1F12/TMPRSS2 polynucleotides are well known in the art and may be employed in the practice of this aspect of the invention.
- 20 In one embodiment, a method for detecting a 20P1F12/TMPRSS2 mRNA in a biological sample comprises producing cDNA from the sample by reverse transcription using at least one primer; amplifying the cDNA so produced using a 20P1F12/TMPRSS2 polynucleotides as sense and antisense primers to amplify 20P1F12/TMPRSS2 cDNAs therein; and detecting the presence of the amplified 20P1F12/TMPRSS2 cDNA. In 25 another embodiment, a method of detecting a 20P1F12/TMPRSS2 gene in a biological sample comprises first isolating genomic DNA from the sample; amplifying the isolated genomic DNA using 20P1F12/TMPRSS2 polynucleotides as sense and antisense primers to amplify the 20P1F12/TMPRSS2 gene therein; and detecting the presence of the amplified 20P1F12/TMPRSS2 gene. Any number of appropriate sense and antisense probe combinations may be designed from the nucleotide sequence 30 provided for 20P1F12/TMPRSS2 (FIG. 1; SEQ ID NO. XX) and used for this purpose.
- In another embodiment, a method of detecting the presence of a 20P1F12/TMPRSS2 protein in a biological sample comprises first contacting the sample with a 20P1F12/TMPRSS2 antibody, a 20P1F12/TMPRSS2-reactive fragment thereof, or a 35 recombinant protein containing an antigen binding region of a 20P1F12/TMPRSS2 antibody; and then detecting the binding of 20P1F12/TMPRSS2 protein in the sample thereto.

Methods for identifying a cell which expresses 20P1F12/TMPRSS2 are also provided. In one embodiment, an assay for identifying a cell which expresses a 20P1F12/TMPRSS2 gene comprises detecting the presence of 20P1F12/TMPRSS2 mRNA in the cell. Methods for the detection of particular mRNAs in cells are well known and include, for example, hybridization assays using complementary DNA probes (such as in situ hybridization using labeled 20P1F12/TMPRSS2 riboprobes, Northern blot and related techniques) and various nucleic acid amplification assays (such as RT-PCR using complementary primers specific for 20P1F12/TMPRSS2, and other amplification type detection methods, such as, for example, branched DNA, SISBA, TMA and the like). Alternatively, an assay for identifying a cell which expresses a 20P1F12/TMPRSS2 gene comprises detecting the presence of 20P1F12/TMPRSS2 protein in the cell or secreted by the cell. Various methods for the detection of proteins are well known in the art and may be employed for the detection of 20P1F12/TMPRSS2 proteins and 20P1F12/TMPRSS2 expressing cells.

Determining the status of 20P1F12/TMPRSS2 expression patterns in an individual may be used to diagnose cancer and may provide prognostic information useful in defining appropriate therapeutic options. Similarly, the expression status of 20P1F12/TMPRSS2 may provide information useful for predicting susceptibility to particular disease stages, progression, and/or tumor aggressiveness. Therefore, another aspect of the invention provides methods and assays for determining 20P1F12/TMPRSS2 expression status and diagnosing cancers which express 20P1F12/TMPRSS2.

In one embodiment, an assay useful in determining the presence of cancer in an individual comprises detecting a significant increase in 20P1F12/TMPRSS2 mRNA or protein expression in a test cell or tissue sample relative to expression levels in the corresponding normal cell or tissue. The presence of 20P1F12/TMPRSS2 mRNA in a colon sample, for example, may indicate the emergence, presence and/or severity of colon cancer, since normal colon does not express 20P1F12/TMPRSS2. In a related embodiment, 20P1F12/TMPRSS2 expression status may be determined at the protein level rather than at the nucleic acid level. For example, such a method or assay would comprise determining the level of 20P1F12/TMPRSS2 protein expressed by cells in a test tissue sample and comparing the level so determined to the level of 20P1F12/TMPRSS2 expressed in a corresponding normal sample. The presence of 20P1F12/TMPRSS2 protein may be evaluated, for example, using immunohistochemical methods. 20P1F12/TMPRSS2 antibodies or binding partners capable of detecting 20P1F12/TMPRSS2 protein expression may be used in a variety of assay formats well known in the art for this purpose.

Peripheral blood may be conveniently assayed for the presence of prostate or colon cancer cells, using RT-PCR to detect 20P1F12/TMPRSS2 expression therein. The presence of RT-PCR amplifiable 20P1F12/TMPRSS2 mRNA may indicate the presence of one of these cancers. RT-PCR detection assays for tumor cells in peripheral blood are currently being evaluated for use in the diagnosis and management of a number of human solid tumors. In the prostate cancer field, these include RT-PCR assays for the detection of cells expressing PSA and PSM (Verkaik et al., 1997, Urol. Res. 25: 373-384; Ghossein et al., 1995, J. Clin. Oncol. 13: 1195-2000; Heston et al., 1995, Clin. Chem. 41: 1687-1688). RT-PCR assays are well known in the art.

In another approach, a recently described sensitive assay for detecting and characterizing carcinoma cells in blood may be used (Racila et al., 1998, Proc. Natl. Acad. Sci. USA 95: 4589-4594). This assay combines immunomagnetic enrichment with multiparameter flow cytometric and immunohistochemical analyses, and is highly sensitive for the detection of cancer cells in blood, reportedly capable of detecting one epithelial cell in 1 ml of peripheral blood.

Methods for detecting and quantifying the expression of 20P1F12/TMPRSS2 mRNA or protein are described herein and use standard nucleic acid and protein detection and quantification technologies well known in the art. Standard methods for the detection and quantification of 20P1F12/TMPRSS2 mRNA include in situ hybridization using labeled 20P1F12/TMPRSS2 riboprobes, Northern blot and related techniques using 20P1F12/TMPRSS2 polynucleotide probes, RT-PCR analysis using primers specific for 20P1F12/TMPRSS2, and other amplification type detection methods, such as, for example, branched DNA, SISBA, TMA and the like. In a specific embodiment, semi-quantitative RT-PCR may be used to detect and quantify 20P1F12/TMPRSS2 mRNA expression as described in the Examples which follow. Any number of primers capable of amplifying 20P1F12/TMPRSS2 may be used for this purpose, including but not limited to the various primer sets specifically described herein. Standard methods for the detection and quantification of protein may be used for this purpose. In a specific embodiment, polyclonal or monoclonal antibodies specifically reactive with the 20P1F12/TMPRSS2 protein may be used in an immunohistochemical assay of biopsied tissue.

The invention further provides kits for the diagnostic and therapeutic applications described or suggested above. Such kits may comprise a carrier means being compartmentalized to receive in close confinement one or more container means such as vials, tubes, and the like, each of the container means comprising one of the separate elements to be used in the method. For example, one of the container means

may comprise a probe which is or can be detectably labeled. Such probe may be an antibody or polynucleotide specific for 20P1F12/TMPRSS2 protein or gene/mRNA, respectively. Where the kit utilizes nucleic acid hybridization to detect the target nucleic acid, the kit may also have containers containing nucleotide(s) for amplification of the target nucleic acid sequence and/or a container comprising a reporter-means, such as a biotin-binding protein, such as avidin or streptavidin, bound to a reporter molecule, such as an enzymatic, florescent, or radionucleotide label.

10

### EXAMPLES

#### EXAMPLE 1: ISOLATION OF cDNA CORRESPONDING TO 20P1F12/TMPRSS2 GENE BY SSH HYBRIDIZATION CLONING AND EXPRESSION ANALYSIS

15

### MATERIALS AND METHODS

#### Cell lines and Human Tissues

20 All human cancer cell lines used in this study were obtained from the ATCC. All cell lines were maintained in DMEM with 10% fetal calf serum. PrEC (primary prostate epithelial cells) were obtained from Clonetics and were grown in PrEBM media supplemented with growth factors (Clonetics).

25 All human prostate cancer xenografts were originally provided by Charles Sawyers (UCLA) (Klein et al., 1997). LAPC-4 AD and LAPC-9 AD xenografts were routinely passaged as small tissue chunks in recipient SCID males. LAPC-4 AI and LAPC-9 AI xenografts were derived as described previously (Klein et al., 1997) and were passaged in castrated males or in female SCID mice.

30

Human tissues for RNA and protein analyses were obtained from the Human Tissue Resource Center (HTRC) at the UCLA (Los Angeles, CA) and from QualTek, Inc. (Santa Barbara, CA). A benign prostatic hyperplasia tissue sample was patient-derived.

35

#### RNA Isolation:

Tumor tissue and cell lines were homogenized in Trizol reagent (Life Technologies, Gibco BRL) using 10 ml/ g tissue or 10 ml/ 10<sup>8</sup> cells to isolate total RNA. Poly A RNA was purified from total RNA using Qiagen's Oligotex mRNA Mini and Midi kits. Total

and mRNA were quantified by spectrophotometric analysis (O.D. 260/280 nm) and analyzed by gel electrophoresis.

Oligonucleotides:

5 The following HPLC purified oligonucleotides were used.

RSACDN (cDNA synthesis primer):

5'TTTTGTACAAGCTT<sub>30</sub>3'

10 Adaptor 1:

5'CTAATACGACTCACTATAGGGCTCGAGCGGCCGCGCCGGGCAGGT3'  
3'GGCCCGTCCA5'

Adaptor 2:

15 5'GTAATACGACTCACTATAGGGCAGCGTGGTTCGCGGCCGAGGT3'  
3'CGGCTCCA5'

PCR primer 1:

5'CTAATACGACTCACTATAGGGC3'

20 Nested primer (NP)1:

5'TCGAGCGGCCGCGCCGGGCAGGT3'

Nested primer (NP)2:

5'AGCGTGGTTCGCGGCCGAGGT3'

25

Suppression Subtractive Hybridization:

Suppression Subtractive Hybridization (SSH) was used to identify cDNAs corresponding to genes which may be up-regulated in androgen dependent prostate cancer compared to benign prostatic hyperplasia.

30

Double stranded cDNAs corresponding to the LAPC-4 AD xenograft (tester) and the BPH tissue (driver) were synthesized from 2 µg of poly(A)<sup>+</sup> RNA isolated from xenograft and BPH tissue, as described above, using CLONTECH's PCR-Select cDNA Subtraction Kit and 1 ng of oligonucleotide RSACDN as primer. First- and second-strand synthesis were carried out as described in the Kit's user manual protocol (CLONTECH Protocol No. PT1117-1, Catalog No. K1804-1). The resulting cDNA was digested with Rsa I for 3 hrs. at 37°C. Digested cDNA was extracted with phenol/chloroform (1:1) and ethanol precipitated.

35

Driver cDNA (BPH) was generated by combining in a 4 to 1 ratio Rsa I digested BPH cDNA with digested cDNA from mouse liver, in order to ensure that murine genes were subtracted from the tester cDNA (LAPC-4 AD).

- 5 Tester cDNA (LAPC-4 AD) was generated by diluting 1  $\mu$ l of Rsa I digested LAPC-4 AD cDNA (400 ng) in 5  $\mu$ l of water. The diluted cDNA (2  $\mu$ l, 160 ng) was then ligated to 2  $\mu$ l of adaptor 1 and adaptor 2 (10  $\mu$ M), in separate ligation reactions, in a total volume of 10  $\mu$ l at 16°C overnight, using 400 u of T4 DNA ligase (CLONTECH). Ligation was terminated with 1  $\mu$ l of 0.2 M EDTA and heating at 72°C for 5 min.

10

- The first hybridization was performed by adding 1.5  $\mu$ l (600 ng) of driver cDNA to each of two tubes containing 1.5  $\mu$ l (20 ng) adaptor 1- and adaptor 2- ligated tester cDNA. In a final volume of 4  $\mu$ l, the samples were overlayed with mineral oil, denatured in an MJ Research thermal cycler at 98°C for 1.5 minutes, and then were allowed to hybridize for 8 hrs at 68°C. The two hybridizations were then mixed together with an additional 1  $\mu$ l of fresh denatured driver cDNA and were allowed to hybridize overnight at 68°C. The second hybridization was then diluted in 200  $\mu$ l of 20 mM Hepes, pH 8.3, 50 mM NaCl, 0.2 mM EDTA, heated at 70°C for 7 min. and stored at -20°C.

20 PCR Amplification, Cloning and Sequencing of Gene Fragments Generated from SSH:

- To amplify gene fragments resulting from SSH reactions, two PCR amplifications were performed. In the primary PCR reaction 1  $\mu$ l of the diluted final hybridization mix was added to 1  $\mu$ l of PCR primer 1 (10  $\mu$ M), 0.5  $\mu$ l dNTP mix (10  $\mu$ M), 2.5  $\mu$ l 10 x reaction buffer (CLONTECH) and 0.5  $\mu$ l 50 x Advantage cDNA polymerase Mix (CLONTECH) in a final volume of 25  $\mu$ l. PCR 1 was conducted using the following conditions: 75°C for 5 min., 94°C for 25 sec., then 27 cycles of 94°C for 10 sec, 66°C for 30 sec, 72°C for 1.5 min. Five separate primary PCR reactions were performed for each experiment. The products were pooled and diluted 1:10 with water. For the secondary PCR reaction, 1  $\mu$ l from the pooled and diluted primary PCR reaction was added to the same reaction mix as used for PCR 1, except that primers NP1 and NP2 (10  $\mu$ M) were used instead of PCR primer 1. PCR 2 was performed using 10-12 cycles of 94°C for 10 sec, 68°C for 30 sec, 72°C for 1.5 minutes. The PCR products were analyzed using 2% agarose gel electrophoresis.

- 35 The PCR products were inserted into pCR2.1 using the T/A vector cloning kit (Invitrogen). Transformed E. coli were subjected to blue/white and ampicillin selection. White colonies were picked and arrayed into 96 well plates and were grown in liquid culture overnight. To identify inserts, PCR amplification was performed on 1 ml of

bacterial culture using the conditions of PCR1 and NP1 and NP2 as primers. PCR products were analyzed using 2% agarose gel electrophoresis.

5 Bacterial clones were stored in 20% glycerol in a 96 well format. Plasmid DNA was prepared, sequenced, and subjected to nucleic acid homology searches of the GenBank, dBest, and NCI-CGAP databases.

**RT-PCR Expression Analysis:**

10 First strand cDNAs were generated from 1 µg of mRNA with oligo (dT)12-18 priming using the Gibco-BRL Superscript Preamplification system. The manufacturers protocol was used and included an incubation for 50 min at 42°C with reverse transcriptase followed by RNase H treatment at 37°C for 20 min. After completing the reaction, the volume was increased to 200 µl with water prior to normalization. First strand cDNAs from 16 different normal human tissues were obtained from Clontech.

15 Normalization of the first strand cDNAs from multiple tissues was performed by using the primers 5'atatgccgcgctcgtcgtcgacaa3' and 5'agccacacgcagctcattgtagaagg 3' to amplify β-actin. First strand cDNA (5 µl) was amplified in a total volume of 50 µl containing 0.4 µM primers, 0.2 µM each dNTPs, 1XPCR buffer (Clontech, 10 mM Tris-HCL, 1.5 mM MgCl<sub>2</sub>, 50 mM KCl, pH8.3) and 1X KlenTaq DNA polymerase (Clontech). Five µl of the PCR reaction was removed at 18, 20, and 22 cycles and used for agarose gel electrophoresis. PCR was performed using an MJ Research thermal  
20 cycler under the following conditions: initial denaturation was at 94°C for 15 sec, followed by a 18, 20, and 22 cycles of 94°C for 15, 65°C for 2 min, 72°C for 5 sec. A  
25 final extension at 72°C was carried out for 2 min. After agarose gel electrophoresis, the band intensities of the 283 bp β-actin bands from multiple tissues were compared by visual inspection. Dilution factors for the first strand cDNAs were calculated to result in equal β-actin band intensities in all tissues after 22 cycles of PCR. Three  
30 rounds of normalization were required to achieve equal band intensities in all tissues after 22 cycles of PCR.

To determine expression levels of the 20P1F12 gene, 5 µl of normalized first strand cDNA was analyzed by PCR using 25, 30, and 35 cycles of amplification using the following primer pairs, which were designed with the assistance of (MIT; for details,  
35 see, [www.genome.wi.mit.edu](http://www.genome.wi.mit.edu)):

5' AGT CTT CCT GCT GAG TCC TTT CC 3'  
5' CAA GGG CAC TGT CTA TAT TCT CAC C 3'



Semi quantitative expression analysis was achieved by comparing the PCR products at cycle numbers that give light band intensities.

5     **RESULTS:**

Several SSH experiments were conducted as described in the Materials and Methods, supra, and led to the isolation of numerous candidate gene fragment clones. All candidate clones were sequenced and subjected to homology analysis against all sequences in the major public gene and EST databases in order to provide information  
10    on the identity of the corresponding gene and to help guide the decision to analyze a particular gene for differential expression.

One of the cDNA clones, designated 20P1F12, showed identity to a recently described serine protease TMPRSS2 (Paoloni-Giacobino et al., 1997, Genomics 44: 309-320).  
15    The isolated 20P1F12 cDNA fragment is 388 bp in length and has the nucleotide sequence shown in FIG. 4. Differential expression analysis by RT-PCR showed that the 20P1F12 gene is expressed at approximately equal levels in normal prostate and the LAPC-4 and LAPC-9 xenografts (FIG. 5, panel A). Further RT-PCR expression analysis of first strand cDNAs from 16 normal tissues showed greatest levels of  
20    20P1F12 expression in prostate. Substantially lower level expression was observed in several other normal tissues (i.e., colon, pancreas, kidney, liver and lung) (FIG. 5, panels B and C).

25    **EXAMPLE 2: NORTHERN BLOT ANALYSIS OF 20P1F12/TMPRSS2 GENE EXPRESSION**

Northern blot analysis on a panel of 16 normal human tissues using a labeled 20P1F12/TMPRSS2 probe (corresponding to the 20P1F12 SSH cDNA of FIG. 4) were conducted to confirm the prostate specificity of 20P1F12/TMPRSS2  
30    expression initially established by RT-PCR expression analysis. The results, shown in FIG. 6 (Panels A & B), confirm and extend the RT-PCR analyses and show that 20P1F12/TMPRSS2 expression is relatively prostate specific, as expression in prostate is clearly many times greater than expression in lung, kidney, pancreas or colon, where only very low level expression is detected.  
35    No detectable expression was observed in any of the other 11 normal tissues used in this panel.

In addition, 20P1F12/TMPRSS2 expression levels in the LAPC-4 and LAPC-9 xenografts were also examined by Northern blot analysis. The results, shown

in FIG. 6 (Panel C), indicate similar expression levels in the xenografts and normal tissue, with lower level expression seen in the LAPC-9 xenograft only. Further Northern blot analysis of 20P1F12/TMPRSS2 expression in a large panel of cancer cells is described in Example 4, below.

5

#### **EXAMPLE 3: CLONING OF FULL LENGTH 20P1F12 cDNA**

A full length cDNA encoding the 20P1F12/TMPRSS2 gene was isolated from a human prostate library and designated 20P1F12-GTC1. The nucleotide and amino acid sequences of 20P1F12-GTC1 are shown in FIG 1. Plasmid p20P1F12-GTC1 (carrying the 20P1F12-GTC1 cDNA) was deposited with the ATCC (Manassas, Virginia) on February 12, 1999 and has been accorded ATCC Designation Number 207097. The approximately 3.5 kb 20P1F12-GTC1 cDNA encodes a protein of 492 amino acids which is almost, but not completely, identical to the sequence previously described (FIG. 2). There are several differences in the nucleotide sequence of the 20P1F12-GTC1 cDNA relative to the published TMPRSS2 sequence, five of which result in different encoded amino acids, as shown in the amino acid alignment of FIG. 3. Specifically, four of the amino acid differences are in the protease domain, three of which are non-conservative amino acid differences which could affect protease function and/or specificity. It is unclear how these amino acid sequence differences might affect biological activity. However, it is possible that 20P1F12/TMPRSS2 and TMPRSS2 are differentially expressed in view of applicants' data showing divergent mRNA expression pattern in normal human tissues.

#### **EXAMPLE 4: 20P1F12/TMPRSS2 EXPRESSION IN PROSTATE AND COLON CANCER**

To analyze 20P1F12/TMPRSS2 expression in cancer tissues and cell lines, Northern blotting was performed on RNA derived from the LAPC xenografts and a panel of prostate and non-prostate cancer cell lines. The results show high levels of 20P1F12/TMPRSS2 expression in all the LAPC xenografts and in colon cancer cell lines (FIG. 7). Similar expression levels were detected in prostate, LAPC-4 AD, LAPC-4 AI, LAPC-9 AD and LNCaP. Lower levels of 20P1F12/TMPRSS2 were seen in LAPC-9 AI and PC-3 cells with no expression seen in DU145. High levels of 20P1F12/TMPRSS2 expression were also detected in three out of four colon cancer cell lines, including LoVo, T84 and Colo-205.

**EXAMPLE 5: CHARACTERIZATION OF 20P1F12/TMPRSS2 PROTEIN****Generation of 20P1F12/TMPRSS2 Monoclonal Antibodies**

TMPRSS2 represents a potential therapeutic target for prostate and colon cancers.

5 As a cell surface antigen, it may be a particularly good target for antibody therapy. To explore this possibility and to further characterize the 20P1F12/TMPRSS2 protein, monoclonal antibodies directed against a GST-20P1F12/TMPRSS2 fusion protein were generated. The immunogen comprised an approximately 8 kD region within the protease domain, specifically amino acid residues 362 through 440 (see FIG. 1). Mice  
10 were immunized with purified GST-TMPRSS2 and hybridomas were generated. Hybridoma supernatants were screened for specific antibodies by western blotting using lysates from 293T cells transfected with 20P1F12/TMPRSS2. A total of 6 hybridomas were identified that specifically recognize 20P1F12/TMPRSS2 by Western blotting (FIG. 8a).

15 Western blotting of LNCaP, LAPC-4 and LAPC-9 cell lysates identifies two major protein bands of approximately 70 and 32 kilodaltons (kD) (FIG. 8b). The predicted molecular weight (MW) of 20P1F12/TMPRSS2 is 54 kD, suggesting that the 70 kD isoform is modified, possibly by glycosylation. The 32 kD form may be a proteolytically  
20 cleaved fragment containing the carboxyl-terminal epitopes recognized by the antibodies.

Additional 20P1F12/TMPRSS2 mAbs may be generated by cell-based immunization using LAPC-9 cells and PC-3 cells expressing 20P1F12/TMPRSS2 as a screening agent  
25 for cell-based ELISAs. In addition, 20P1F12/TMPRSS2 mAbs may be generated using purified 20P1F12/TMPRSS2 protein as immunogen. For example, recombinant 20P1F12/TMPRSS2 having an amino-terminal His-tag may be expressed in a baculovirus system using pBlueBac4.5 (Invitrogen). His-tagged 20P1F12/TMPRSS2 may then be purified using a Nickel column, quantified and used as immunogen.  
30 Screening of monoclonal may be performed using cell-based ELISAs with, for example, LNCaP and PC-3/TMPRSS2 cells.

**Cell Surface Localization**

To study the characteristics of the 20P1F12/TMPRSS2 protein, 20P1F12 cDNA (FIG. 1)  
35 was cloned into pcDNA 3.1 Myc-His (Invitrogen), which provides a 6-His tag at the carboxyl-terminus. The construct was transfected into 293T cells and was analyzed by cell-surface biotinylation. Biotinylated cell surface proteins were affinity purified using streptavidin-sepharose. Western blot analysis of streptavidin affinity purified proteins using an anti-His antibody demonstrated the presence of 20P1F12/TMPRSS2

protein (FIG. 9a). Therefore, as predicted from sequence analysis, 20P1F12/TMPRSS2 is expressed at the cell surface of transfected cells.

5 To examine cell surface expression of endogenous 20P1F12/TMPRSS2 in LNCaP and PC-3 prostate cancer cells, biotinylated cell surface proteins were affinity purified with streptavidin-sepharose and probed with anti-20P1F12/TMPRSS2 antibodies. Western blotting of streptavidin purified proteins clearly show cell surface biotinylation of endogenous 20P1F12/TMPRSS2 in both LNCaP and PC-3 cells appearing as 32 and 70 kD protein bands (FIG. 9b). In additional controls, 10 20P1F12/TMPRSS2 protein was not detected in streptavidin precipitates from non-biotinylated cells (Fig. 8b). This data combined with sequence analysis predict 20P1F12/TMPRSS2 to be a type II transmembrane protein.

15 Interestingly, 293T cells transfected with a carboxyl-terminal His-tagged 20P1F12/TMPRSS2 express primarily the 70 kD protein (FIG. 9a). Since the 20P1F12/TMPRSS2 protease domain is located at the carboxyl-terminus, it is possible that the 32 kD fragment is a result of auto-catalytic cleavage, which is inhibited by the His tag. The related molecule, hepsin (TMPRSS1), appears to be capable of autoactivation in a concentration dependent manner (Vu et al., 1997, J. Biol. Chem. 20 272: 31315-31320). This auto-catalytic cleavage may be exploited to identify small molecules that inhibit 20P1F12/TMPRSS2 activity. Cells may be grown in the presence or absence of small molecule inhibitors to specifically look for inhibition of cleavage. Such small molecules may be tested as prostate cancer therapeutics.

#### 25 Glycosylation of 20P1F12/TMPRSS2

The predicted MW of 20P1F12/TMPRSS2 is significantly smaller than the apparent MW detected by Western blotting. This suggests that 20P1F12/TMPRSS2 may be glycosylated. The GTC1 sequence indicates that there are three potential glycosylation sites with the consensus sequence of NXS/T (residues 128, 213, 249). To 30 explore the possibility that 20P1F12/TMPRSS2 is glycosylated, His-tagged 20P1F12/TMPRSS2 was transfected into 293T cells and purified using a Nickel-agarose (Invitrogen). Affinity purified protein was eluted with 50 mM EDTA, pH 8.0, and was de-glycosylated using N-glycosidase F (Boehringer Mannheim) according to the manufacturers protocol. Untreated and de-glycosylated protein were analyzed by 35 western blotting using anti-His antibodies. The results show a 5-8 kD MW shift of 20P1F12/TMPRSS2 with N-glycosidase F treatment (FIG 10), indicating that 20P1F12/TMPRSS2 is indeed a glycosylated protein. De-glycosylated 20P1F12/TMPRSS2 still exhibited a MW of at least 5-10 kD larger than the predicted size, indicating that either the de-glycosylation reaction was not complete (or that

glycosylation is O-linked), or that 20P1F12/TMPRSS2 may exhibit additional post-translational modifications (such as phosphorylation, sulfation).

#### Androgen Regulation

- 5 Northern blotting shows that expression of 20P1F12/TMPRSS2 seems to decrease in the androgen independent LAPC-9 xenograft and the androgen independent cell lines PC-3 and DU145 (FIG. 6), suggesting that 20P1F12/TMPRSS2 may be an androgen regulated gene. To explore this possibility, LNCaP cells, which are androgen dependent and express significant levels of 20P1F12/TMPRSS2, were deprived of
- 10 androgen for one week by growing them in media containing 2% charcoal-stripped fetal bovine serum (FBS). The cells were then stimulated with mibolerone, a synthetic androgen analogue, at various time points. Cells were harvested for RNA and Northern blotting. As a loading control, the same blot was also probed with  $\beta$ -actin. The results (FIG. 11) show a clear reduction of 20P1F12/TMPRSS2 expression during
- 15 androgen deprivation (FIG. 11). Addition of mibolerone increased 20P1F12/TMPRSS2 expression significantly, indicating that it is an androgen responsive gene. Expression of prostate-specific antigen (PSA) in the same samples was monitored as a positive control for androgen regulation (FIG. 11).
- 20 To determine the optimal time of 20P1F12/TMPRSS2 induction, androgen starved cells were stimulated with mibolerone for various time points. Cells were harvested for RNA and protein isolation to perform northern and western blotting respectively. The results (FIG. 12) show induction of 20P1F12/TMPRSS2 message within three hours of stimulation and increased through 24 hours after hormone addition.
- 25 To analyze the protein levels, western blotting of cell lysates using the 1F9 mAb was performed. Additional controls for 20P1F12/TMPRSS2 expression included PC-3 cells infected with a retrovirus encoding either neo or 20P1F12/TMPRSS2. Infected PC-3 cells were selected in G418 for 2-3 weeks and harvested for western blotting. The
- 30 results showed strong expression of 20P1F12/TMPRSS2 in the cells infected with a 20P1F12/TMPRSS2 virus, and no detectable 20P1F12/TMPRSS2 expression in the neo cells.
- 35 When looking at androgen deprived LNCaP cells, 20P1F12/TMPRSS2 expression is still detectable, but visibly reduced when compared to androgen stimulated cells. However, the first time point of induced expression appears after 9 hours of stimulation, indicating that protein expression of 20P1F12/TMPRSS2 lags behind RNA induction (FIG. 12).

These results demonstrate that 20P1F12/TMPRSS2 is an androgen regulated gene, similar to other prostate specific proteases, such as PSA and hK2 (Young et al., 1995, J. Androl. 16:97).

5     Effect of 20P1F12/TMPRSS2 on NIH 3T3 Morphology

20P1F12/TMPRSS2 exhibits prostate specific expression and seems to be regulated by androgen. To determine the effect of expressing 20P1F12/TMPRSS2 in a heterologous non-prostate cancer cell line, 20P1F12/TMPRSS2 retrovirus was used to infect NIH 3T3 cells. The morphology of cells infected with 20P1F12/TMPRSS2  
10     retrovirus was compared to the morphology of control (neo) virus infected cells. A population of infected cells exhibited a distinct vacuolar appearance compared to control cells (FIG. 13), which seem to correlate with high levels of expression. Upon passaging this infected cell population, vacuole-bearing cells gradually disappeared with apparently reduced expression of 20P1F12/TMPRSS2.

15     Evaluation of 20P1F12/TMPRSS2 Function

20P1F12/TMPRSS2 function may be being assessed in mammalian cells engineered to express 20P1F12/TMPRSS2. For this purpose, 20P1F12/TMPRSS2 is conveniently cloned into several vectors, including pcDNA 3.1 myc-His-tag (Invitrogen), the  
20     retroviral vector pSR $\alpha$ tkneo (Muller et al., 1991, MCB 11:1785), and pIND (Invitrogen) an ecdysone-inducible expression system. Using these expression vectors, 20P1F12/TMPRSS2 is expressed in several cell lines, including PC-3, NIH 3T3, mouse L cell fibroblasts and 293T. Expression of 20P1F12/TMPRSS2 is monitored using anti-20P1F12/TMPRSS2 antibodies by Western and FACS analysis. Purified TMPRSS2 may  
25     used to identify the substrate.

Such mammalian cell lines expressing 20P1F12/TMPRSS2 are then tested in several in vitro, including cell proliferation, cell adhesion, cell invasion using a membrane invasion culture system (MICS) (Welch et al., Int. J. Cancer 43: 449-457) in tissue  
30     culture, and in vivo assays, including tumor formation in SCID mice. The 20P1F12/TMPRSS2 cell phenotype is compared to the phenotype of cells which do not express 20P1F12/TMPRSS2.

To assess the functional role of the different domains in 20P1F12/TMPRSS2, the  
35     following deletion mutants and point mutants are generated: (i)  $\Delta$ SRCR (93 a.a. deletion); (ii)  $\Delta$ LDLRA (35 a.a. deletion); and (iii) mutant of the catalytic triad: H296Q, D345N, S441A (single point mutants). 20P1F12/TMPRSS2 mutants are cloned into the retroviral vectors pSR $\alpha$ tkneo for expression in mammalian cells. The resulting mutants are useful for elucidating the importance of the different domains and

residues. In addition, these experiments are useful for determining whether such mutants function as dominant negative molecules. Dominant negative activity may be manifested in cells that express endogenous 20P1F12/TMPRSS2, such as LNCaP. Dominant negative activity may be due to interactions with substrates via protease domain, or via the protein-protein interaction domains. The mutant 20P1F12/TMPRSS2 molecules are tested in the same in vitro and in vivo assays as wild-type 20P1F12/TMPRSS2 (see above). Such dominant negative 20P1F12/TMPRSS2 molecules may be useful therapeutically. For example, a dominant negative 20P1F12/TMPRSS2 may introduced into prostate cancer cells via gene therapy vectors capable of delivering and expressing the corresponding coding sequence into prostate tumor cells. Similarly, such methods may be useful in the treatment of colon cancer.

Determining the characteristics of 20P1F12/TMPRSS2 expression in normal mouse tissues and in transgenic mice provides further information about the function of 20P1F12/TMPRSS2. Northern blot analysis using probes designed from the 20P1F12/TMPRSS2 sequences provided herein may be used to define the expression pattern of murine 20P1F12/TMPRSS2. In addition, 20P1F12/TMPRSS2 expression during development in the mouse embryo can be analyzed. The resulting data will identify a tissue source for cloning the mouse gene and predict which tissues would be affected in a transgenic mouse knock-out study.

transgenic mouse may be generated and used to define the biological role of 20P1F12/TMPRSS2 in an in-vivo setting. In one approach, the human or mouse 20P1F12/TMPRSS2 genes are used to generate transgenic mice. Over-expression of spontaneous tumor formation in mice may be studies using transgenic mice. In another approach, 20P1F12/TMPRSS2 gene knock-outs are generated in mice. Such mice may also be crossed with other prostate cancer mouse models, such as the TRAMP model (Greenberg et al., 1995, PNAS 92:3439) to study the influence on prostate cancer aggressiveness and metastasis and to observe changes in disease progression.

Experiments testing 20P1F12/TMPRSS2 functional interaction with serine protease inhibitors will also provide information on 20P1F12/TMPRSS2 function. For this purpose, inhibition is accomplished using small molecule inhibitors or biological inhibitors.

- 5 Throughout this application, various publications are referenced within parentheses. The disclosures of these publications are hereby incorporated by reference herein in their entireties.

- 10 The present invention is not to be limited in scope by the embodiments disclosed herein, which are intended as single illustrations of individual aspects of the invention, and any which are functionally equivalent are within the scope of the invention. Various modifications to the models and methods of the invention, in addition to those described herein, will become apparent to those skilled in the art from the foregoing description and teachings, and are similarly intended to fall within the scope of the
- 15 invention. Such modifications or other embodiments can be practiced without departing from the true scope and spirit of the invention.



**CLAIMS:**

1. An isolated 20P1F12/TMPRSS2 protein having an amino acid sequence shown in FIG. 1 (SEQ ID NO. XX).
- 5 2. An isolated polynucleotide selected from the group consisting of (a) a polynucleotide having the sequence as shown in FIG. 1 (SEQ ID NO. XX), wherein T can also be U; (b) a polynucleotide encoding a 20P1F12/TMPRSS2 polypeptide whose sequence is encoded by the cDNA contained in plasmid p20P1F12-GTC1  
10 as deposited with American Type Culture Collection as Accession No. 207097; and (c) a polynucleotide encoding the 20P1F12/TMPRSS2 protein of claim 1.
3. An isolated polynucleotide which is fully complementary to a polynucleotide according to claim 2.
- 15 4. A recombinant expression vector which contains a polynucleotide according to claim 3.
5. A host cell which contains an expression vector according to claim 4.
- 20 6. An antibody which immunospecifically binds to the 20P1F12/TMPRSS2 protein of claim 1.
7. A monoclonal antibody according to claim 6.
- 25 8. A fragment of the antibody of claim 7.
9. A recombinant protein comprising the antigen binding domain of the antibody of claim 7.
- 30 10. The antibody of claim 7 which is labeled with a detectable marker.
11. The monoclonal antibody of claim 7 which is conjugated to a toxin.
- 35 12. The monoclonal antibody of claim 7 which is conjugated to a therapeutic agent.
13. The antibody fragment of claim 8 which is labeled with a detectable marker.
14. The recombinant protein of claim 9 which is labeled with a detectable marker.

15. An assay for detecting the presence of a 20P1F12/TMPRSS2 protein in a biological sample comprising contacting the sample with an antibody of claim 10, 13 or 14 and detecting the binding of 20P1F12/TMPRSS2 protein in the sample thereto.
16. An assay for detecting the presence of a 20P1F12/TMPRSS2 polynucleotide in a biological sample, comprising
- (a) contacting the sample with a polynucleotide probe which specifically hybridizes to the 20P1F12/TMPRSS2 cDNA contained within plasmid p20P1F12-GTC1 as deposited with American Type Culture Collection as Accession No. 207097, or the polynucleotide as shown in FIG. 1 (SEQ ID NO. XX), or the complements thereof; and
- (b) detecting the presence of a hybridization complex formed by the hybridization of the probe with 20P1F12/TMPRSS2 polynucleotide in the sample, wherein the presence of the hybridization complex indicates the presence of 20P1F12/TMPRSS2 polynucleotide within the sample.
17. An assay for detecting the presence of 20P1F12/TMPRSS2 mRNA in a biological sample comprising:
- (a) producing cDNA from the sample by reverse transcription using at least one primer;
- (b) amplifying the cDNA so produced using 20P1F12/TMPRSS2 polynucleotides as sense and antisense primers to amplify 20P1F12/TMPRSS2 cDNAs therein;
- (c) detecting the presence of the amplified 20P1F12/TMPRSS2 cDNA,
- wherein the 20P1F12/TMPRSS2 polynucleotides used as the sense and antisense probes are capable of amplifying the polynucleotide shown in FIG. 1 (SEQ ID NO. XX).
18. A composition for the treatment of prostate cancer comprising an antibody according to claim 7, 11 or 12, wherein the antibody binds to an extracellular domain of 20P1F12/TMPRSS2.

37. A composition for the treatment of colon cancer comprising an antibody according to claim 7, 11 or 12, wherein the antibody binds to an extracellular domain of 20P1F12/TMPRSS2.
- 5

5' GGC GGA GGC GGA GGC GGA GGG CGA GGG GCG GGG AGC GCC GCC TGG AGC GCG GCA  
-----  
GGT CAT ATT GAA CAT TCC AGA TAC CTA TCA TTA CTC GAT GCT GTT GAT AAC AGC  
-----  
AAG ATG GCT TTG AAC TCA GGG TCA CCA CCA GCT ATT GGA CCT TAC TAT GAA AAC  
-----  
M A L N S G S P P A I G P Y Y E N  
-----  
CAT GGA TAC CAA CCG GAA AAC CCC TAT CCC GCA CAG CCC ACT GTG GTC CCC ACT  
-----  
H G Y Q P E N P Y P A Q P T V V P T  
-----  
GTC TAC GAG GTG CAT CCG GCT CAG TAC TAC CCG TCC CCC GTG CCC CAG TAC GCC  
-----  
V Y E V H P A Q Y Y P S P V P Q Y A  
-----  
CCG AGG GTC CTG ACG CAG GCT TCC AAC CCC GTC GTC TGC ACG CAG CCC AAA TCC  
-----  
P R V L T Q A S N P V V C T Q P K S  
-----  
CCA TCC GGG ACA GTG TGC ACC TCA AAG ACT AAG AAA GCA CTG TGC ATC ACC TTG  
-----  
P S G T V C T S K T K K A L C I T L  
-----  
ACC CTG GGG ACC TTC CTC GTG GGA GCT GCG CTG GCC GCT GGC CTA CTC TGG AAG  
-----  
T L G T F L V G A A L A A G L L W K  
-----  
TTC ATG GGC AGC AAG TGC TCC AAC TCT GGG ATA GAG TGC GAC TCC TCA GGT ACC  
-----  
F M G S K C S N S G I E C D S S G T  
-----  
TGC ATC AAC CCC TCT AAC TGG TGT GAT GGC GTG TCA CAC TGC CCC GGC GGG GAG  
-----  
C I N P S N W C D G V S H C P G G E  
-----  
GAC GAG AAT CGG TGT GTT CGC CTC TAC GGA CCA AAC TTC ATC CTT CAG GTG TAC  
-----  
D E N R C V R L Y G P N F I L Q V Y  
-----  
TCA TCT CAG AGG AAG TCC TGG CAC CCT GTG TGC CAA GAC GAC TGG AAC GAG AAC  
-----  
S S Q R K S W H P V C Q D D W N E N  
-----  
TAC GGG CGG GCG GCC TGC AGG GAC ATG GGC TAT AAG AAT AAT TTT TAC TCT AGC  
-----  
Y G R A A C R D M G Y K N N F Y S S

CAA	GGA	ATA	GTG	GAT	GAC	AGC	GGA	TCC	ACC	AGC	TTT	ATG	AAA	CTG	AAC	ACA	AGT
Q	G	I	V	D	D	S	G	S	T	S	F	M	K	L	N	T	S
GCC	GGC	AAT	GTC	GAT	ATC	TAT	AAA	AAA	CTG	TAC	CAC	AGT	GAT	GCC	TGT	TCT	TCA
A	G	N	V	D	I	Y	K	K	L	Y	H	S	D	A	C	S	S
AAA	GCA	GTG	GTT	TCT	TTA	CGC	TGT	ATA	GCC	TGC	GGG	GTC	AAC	TTG	AAC	TCA	AGC
K	A	V	V	S	L	R	C	I	A	C	G	V	N	L	N	S	S
CGC	CAG	AGC	AGG	ATT	GTG	GGC	GGC	GAG	AGC	GCG	CTC	CCG	GGG	GCC	TGG	CCC	TGG
R	Q	S	R	I	V	G	G	E	S	A	L	P	G	A	W	P	W
CAG	GTC	AGC	CTG	CAC	GTC	CAG	AAC	GTC	CAC	GTG	TGC	GGA	GGC	TCC	ATC	ATC	ACC
Q	V	S	L	H	V	Q	N	V	H	V	C	G	G	S	I	I	T
CCC	GAG	TGG	ATC	GTG	ACA	GCC	GCC	CAC	TGC	GTG	GAA	AAA	CCT	CTT	AAC	AAT	CCA
P	E	W	I	V	T	A	A	H	C	V	E	K	P	L	N	N	P
TGG	CAT	TGG	ACG	GCA	TTT	GCG	GGG	ATT	TTG	AGA	CAA	TCT	TTC	ATG	TTC	TAT	GGA
W	H	W	T	A	F	A	G	I	L	R	Q	S	F	M	F	Y	G
GCC	GGA	TAC	CAA	GTA	GAA	AAA	GTG	ATT	TCT	CAT	CCA	AAT	TAT	GAC	TCC	AAG	ACC
A	G	Y	Q	V	E	K	V	I	S	H	P	N	Y	D	S	K	T
AAG	AAC	AAT	GAC	ATT	GCG	CTG	ATG	AAG	CTG	CAG	AAG	CCT	CTG	ACT	TTC	AAC	GAC
K	N	N	D	I	A	L	M	K	L	Q	K	P	L	T	F	N	D
CTA	GTG	AAA	CCA	GTG	TGT	CTG	CCC	AAC	CCA	GGC	ATG	ATG	CTG	CAG	CCA	GAA	CAG
L	V	K	P	V	C	L	P	N	P	G	M	M	L	Q	P	E	Q
CTC	TGC	TGG	ATT	TCC	GGG	TGG	GGG	GCC	ACC	GAG	GAG	AAA	GGG	AAG	ACC	TCA	GAA
L	C	W	I	S	G	W	G	A	T	E	E	K	G	K	T	S	E
GTG	CTG	AAC	GCT	GCC	AAG	GTG	CTT	CTC	ATT	GAG	ACA	CAG	AGA	TGC	AAC	AGC	AGA
V	L	N	A	A	K	V	L	L	I	E	T	Q	R	C	N	S	R
TAT	GTC	TAT	GAC	AAC	CTG	ATC	ACA	CCA	GCC	ATG	ATC	TGT	GCC	GGC	TTC	CTG	CAG
Y	V	Y	D	N	L	I	T	P	A	M	I	C	A	G	F	L	Q

1415	1424	1433	1442	1451	1460
GGG AAC GTC GAT TCT TGC CAG GGT GAC AGT GGA GGG CCT CTG GTC ACT TCG AAG					
-----	-----	-----	-----	-----	-----
G N V D S C Q G D S G G P L V T S K					
1469	1478	1487	1496	1505	1514
AAC AAT ATC TGG TGG CTG ATA GGG GAT ACA AGC TGG GGT TCT GGC TGT GCC AAA					
-----	-----	-----	-----	-----	-----
N N I W W L I G D T S W G S G C A K					
1523	1532	1541	1550	1559	1568
GCT TAC AGA CCA GGA GTG TAC GGG AAT GTG ATG GTA TTC ACG GAC TGG ATT TAT					
-----	-----	-----	-----	-----	-----
A Y R P G V Y G N V M V F T D W I Y					
1577	1586	1595	1604	1613	1622
CGA CAA ATG AGG GCA GAC GGC TAA TCC ACA TGG TCT TCG TCC TTG ACG TCG TTT					
-----	-----	-----	-----	-----	-----
R Q M R A D G *					
1631	1640	1649	1658	1667	1676
TAC AAG AAA ACA ATG GGG CTG GTT TTG CTT CCC CGT GCA TGA TTT ACT CTT AGA					
-----	-----	-----	-----	-----	-----
1685	1694	1703	1712	1721	1730
GAT GAT TCA GAG GTC ACT TCA TTT TTA TTA AAC AGT GAA CTT GTC TGG CAA AAA					
-----	-----	-----	-----	-----	-----
1739					
AAA AAA AAA A 3'					
-----					

1 gtcattattga acattccaga tacctatcat tactcgatgc tgttgataac agcaagatgg  
61 ctttgaactc aggggtcacca ccagctattg gaccttacta tgaaaacccat ggataccaac  
121 cggaaaaccc ctatcccga cagcccaactg tggccccac tgtctacgag gtgcatccgg  
181 ctcagtacta cccgtccccc gtgccccagt acgccccgag ggtcctgacg caggcttcca  
241 acccgcgtgt ctgcacgcag cccaaatccc catccgggac agtgtgcacc tcaaagacta  
301 agaaagcact gtgcatcacc ttgaccctgg ggaccttcct cgtggggagct gcgctggccg  
361 ctggcctact ctggaagtgc atggggcagca agtgcctcaa ctctgggata gagtgcgact  
421 cctcaggtac ctgcatcaac ccccttaact ggtgtgatgg cgtgtcacac tgccccggcg  
481 gggaggacga gaatcgggtg gttcgcctct acggaccaaa cttcatcctt cagatgtact  
541 catctcagag gaagtcctgg caccctgtgt gccaaagcga ctggaacgag aactacgggc  
601 gggcggcctg cagggacatg ggctataaga ataattttta ctctagccaa ggaatagtgg  
661 atgacagcgg atccaccagc tttatgaaac tgaacacaag tgccggcaat tccgatattc  
721 ataaaaaact gtaccacagt gatgcctgtt cttcaaaagc agtgggtttct ttacgctgtt  
781 tagcctgcgg ggtcaacttg aactcaagcc gccagagcag gatcgtgggc ggtgagagcg  
841 cgctcccggg ggcctggccc tggcaggtca gcctgcacgt ccagaacgct cactgtgtgc  
901 gaggtcccat catcaccccc gagtggatcg tgacagccgc ccactgcgtg gaaaaacctc  
961 ttaacaatcc atggcattgg acggcatttg cggggatttt gagacaatct ttcattgtct  
1021 atggagccgg ataccaagta caaaaagtga tttctcatcc aaattatgac tccaagacca  
1081 agaacaatga cattgcgctg atgaagctgc agaagcctct gactttcaac gacctagtga  
1141 aaccagtgtg tctgccccac ccaggcatga tgctgcagcc agaacagctc tgctggattt  
1201 ccgggtgggg ggccaccgag gagaaaggga agacctcaga agtgcagaac gctgccaagg  
1261 tgcttctcat tgagacacag agatgcaaca gcagatatgt ctatgacaac ctgatcacac  
1321 cagccatgat ctgtgccggc ttcttcaggg ggaacgtcga ttcttgccag ggtgacagtg  
1381 gagggcctct ggtcacttcg aacaacaata tctggtggct gataggggat acaagctggg  
1441 gttctggctg tgccaaagct tacagaccag gagtgtacgg gaatgtgatg gtattcacgg  
1501 actggattta tcgacaaatg aaggcaaacg gctaattcac atggctcttcg tccctgacgt  
1561 cgtttttaca gaaaacaatg gggctgggtt tgcttccccg tgcatgattt actcttagag  
1621 atgattcaga ggtcacttca tttttattaa acagtgaact tgtctggctt tggcactctc  
1681 tgccatactg tgcaggctgc agtggctccc ctgcccagcc tgctctccct aacccttgt  
1741 ccgcaagggg tgatggccgg ctggttgtgg gcactggcgg tcaattgtgg aaggaagagg  
1801 gttggaggct gccccattg agatcttctt gctgagtcct ttccagggc caattttgga  
1861 tgagcatgga gctgtcactt ctcagctgct ggatgacttg agatgaaaaa ggagagacat  
1921 ggaaaggagg acagccaggt ggcacctgca gcggctgccc tctggggcca cttggtagtg  
1981 tccccagcct acttcacaag gggattttgc tgatgggttc ttagagcctt agcagccctg  
2041 gatggtggcc agaaataaag ggaccagccc ttcattgggtg gtgacgtggt agtcacttgt  
2101 aaggggaaca gaaacatttt tgttcttatg gggtgagaat atagacagt cccttggtgc  
2161 gaggggaagca attgaaaagg aacttgcctt gagcactcct ggtgcaggtc tccacctgca  
2221 cattgggtgg ggctcctggg agggagactc agccttcctc ctcatcctcc ctgaccctgc  
2281 tcctagcacc ctggagagtg aatgcccctt ggtcccctgg agggcgccaa gtttggcacc  
2341 atgtcggcct cttcaggcct gatagtcatt ggaaattgag gtccatgggg gaaatcaagg  
2401 atgctcagtt taaggtaac tgtttccatg ttatgtttct acacattgat ggtggtgacc  
2461 ctgagttcaa agccatctt

## ORF AMINO ACID SEQUENCE

MALNSGSPPAIGPPYYENHGYQFENPYPAQPTVVPTVYEVHQAQYYPSVPQYAPRVLTQASNPVVCTQPKSPSGTV  
CTSKTKKALCITLTGLTFLVGAALAAGLLWKFMGSKCSNSGIECDSSGTCINPSNWCDGVSHCPGGEDENRCVRLY  
GPNFILQMYSSQRKSWHPVCQDDWNENYGRAACRDMGYKNNFYSSQGI VDDSGSTSFMKLNTSAGNVDIYKKLYHS  
DACSSKAVVSLRCLACGVNLSSRSRIVGGESALPGAWPWQVSLHVQNVHVCSSII TPEWIVTAHCVEKPLNN  
PWHWTAFAFILRQSFMYGAGYQVQKVI SHPNYDSKTKNNDIALMKLQKPLTFNDLVKPVCLPNPGMMLQPEQLCW  
ISGWGATEEKGKTSEVLNAAKVLIIETQRCNSRYVDNLITPAMICAGFLQGNVDSQCQDSSGGLVTSNNNIWWLI  
GDTSWGSGCAKAYRPGVYGNVMVFTDWIYRQMKANG

## FIG 3

1 15 16 30 31 45 46 60 61 75 76 90  
GTC1 MALNSGSPPAIGPY ENHGYQENPYPAQ TVPTVVEVHPAQY PSPVPQYAPRVLTOA SNPVVCTQPKSPSGT VCTSKTKKALCITLT 90  
TMPRSS2 MALNSGSPPAIGPY ENHGYQENPYPAQ TVPTVVEVHPAQY PSPVPQYAPRVLTOA SNPVVCTQPKSPSGT VCTSKTKKALCITLT 90

91 105 106 120 121 135 136 150 151 165 166 180  
GTC1 LGTFLVGAALAGLL WKFMGSKCSNSGIEC DSSGTCINPSNWCDG VSHCPGGEDENRCVR LYGNPFILOQYSSOR KSWHPVCQDDWNENY 180  
TMPRSS2 LGTFLVGAALAGLL WKFMGSKCSNSGIEC DSSGTCINPSNWCDG VSHCPGGEDENRCVR LYGNPFILOQYSSOR KSWHPVCQDDWNENY 180

181 195 196 210 211 225 226 240 241 255 256 270  
GTC1 GRAACRDMGYKNFY SSQGIVDDSGSTSEM KLNTSAGNVDIYKKL YHSDACSSKAVVSLR CIACGVNLNSSRQSR IVGGESALPGAWPWQ 270  
TMPRSS2 GRAACRDMGYKNFY SSQGIVDDSGSTSEM KLNTSAGNVDIYKKL YHSDACSSKAVVSLR CIACGVNLNSSRQSR IVGGESALPGAWPWQ 270

271 285 286 300 301 315 316 330 331 345 346 360  
GTC1 VSLHVQNPHVCGGSI ITPEWIVTAHCVEK PLNNPWHHTAFAGIL RQSFMYGAGYQVQK VISHPNYDSKTKNND IALMKLQKPLTFNDL 360  
TMPRSS2 VSLHVQNPHVCGGSI ITPEWIVTAHCVEK PLNNPWHHTAFAGIL RQSFMYGAGYQVQK VISHPNYDSKTKNND IALMKLQKPLTFNDL 360


361 375 376 390 391 405 406 420 421 435 436 450  
GTC1 VKPVCLPNPGMMLQP EQLCWISGWGATEEK GKTSEVLNAAKVILLI ETORCNSRYVDNLI TPAMICAGFLOQNVND SCQDGSGLPLVTSKN 450  
TMPRSS2 VKPVCLPNPGMMLQP EQLCWISGWGATEEK GKTSEVLNAAKVILLI ETORCNSRYVDNLI TPAMICAGFLOQNVND SCQDGSGLPLVTSNN 450

451 465 466 480 481 495 496 510 511 525 526 540  
GTC1 NIWWLIGDTSWGGC AKAYRPGVYGNVMVF TDWIYRQMRADG 492 540  
TMPRSS2 NIWWLIGDTSWGGC AKAYRPGVYGNVMVF TDWIYRQMKANG 492



## FIG 4

GATCTTCCTGCTGAGTCCTTTCCAGGGGCCAATTTTGGATGAGCATGGAGCTGTCACCTCTCAGCTGCTGGATGAC  
TTGAGATGAAAAGGAGAGACATGGAAAGGGAGACAGCCAGGTGGCACCTGCAGCGGCTGCCCTCTGGGGCCACTT  
GGTAGTGTCCCGAGCCTACCTCTCCACAAGGGGATTTTGCTGATGGGTCTTANAGCCTTAGCAGCCCTGGATGGT  
GGCCAGAAATAAAGGGACCAGCCCTTCATGGGTGGTGACGTGGTANTCACTTGTAAGGGGAACAGAAACATTTTG  
TTCTTATGGGGTGAGAATATAGACAGTGCCCTTGGTGCGAGGGAAGCAATTGAAAAGGAACTTGCCCTGAGCACTC  
CTGGTGCA

**FIG. 5****A**

1. Brain
2. Prostate
3. LAPC-4 AD
4. LAPC-4 AI
5. LAPC-9 AD
6. HeLa
7. Murine cDNA
8. Neg. control

**B**

1. Brain
2. Heart
3. Kidney
4. Liver
5. Lung
6. Pancreas
7. Placenta
8. Skeletal Muscle

**C**

1. Colon
2. Ovary
3. Leukocytes
4. Prostate
5. Small Intestine
6. Spleen
7. Testis
8. Thymus

FIG. 6




FIG. 7







FIG. 8

A




B




**FIG. 9****A. In transfected 293T cells:****B. In prostate cancer cells:**

## FIG. 10




## FIG. 11



## Panel:


1. LNCaP androgen-deprived 1 week
2. LNCaP FBS
3. LNCaP androgen-deprived 24 hrs + mock 9 hrs
4. LNCaP androgen deprived 24 hrs + Mib 9hrs





LNCaP cells were androgen deprived for 1 week (grown in 2% CS-FBS) and were then stimulated with 10 nM mibolerone for various time points

FIG. 13



NIH 3T3  
TMPRSS2

NIH 3T3  
neo



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification 6 :</b> C12N 9/64, 15/57, C07K 14/47, 16/40, C12Q 1/68, A61K 39/395		<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 99/62942</b>
			<b>(43) International Publication Date:</b> 9 December 1999 (09.12.99)
<b>(21) International Application Number:</b> PCT/US99/12253		<b>(81) Designated States:</b> AT, AT (Utility model), AU, BR, CA, CH, CN, DE, DE (Utility model), DK, DK (Utility model), ES, FI, FI (Utility model), GB, IL, JP, KR, MX, NO, NZ, RU, SE, SG, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
<b>(22) International Filing Date:</b> 1 June 1999 (01.06.99)		<b>Published</b> <i>With international search report.</i>	
<b>(30) Priority Data:</b> 60/087,598 1 June 1998 (01.06.98) US 60/091,474 29 June 1998 (29.06.98) US 60/129,521 14 April 1999 (14.04.99) US			
<b>(71) Applicant:</b> UROGENESYS, INC. [US/US]; 1701 Colorado Avenue, Santa Monica, CA 90404 (US).		<b>(88) Date of publication of the international search report:</b> 2 June 2000 (02.06.00)	
<b>(71)(72) Applicants and Inventors:</b> AFAR, Daniel, E. [CA/US]; 17250 Sunset Boulevard, Pacific Palisades, CA 90272 (US). HUBERT, Rene, S. [US/US]; 1503 Westerly Terrace, Los Angeles, CA 90026 (US). LEONG, Kahan [US/US]; 8160 Manitoba Street No. 109, Playa Del Rey, CA 90293 (US). RAITANO, Arthur, B. [US/US]; 10807 Cushdon Avenue, Los Angeles, CA 90064 (US). SAFFRAN, Douglas, C. [-US]; 11720 Bellagio Road No. 204, Los Angeles, CA 90049 (US).			
<b>(74) Agent:</b> SHARPLES, Kenneth, K.; Law Office of Kenneth K. Sharples, Suite 202, 80 Fourth Street, P.O. Box 277, Point Reyes, CA 94956 (US).			

**(54) Title:** TUMOR ANTIGEN USEFUL IN DIAGNOSIS AND THERAPY OF PROSTATE AND COLON CANCER

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080 1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1100 1101 1102 1103 1104 1105 1106 1107 1108 1109 1110 1111 1112 1113 1114 1115 1116 1117 1118 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1140 1141 1142 1143 1144 1145 1146 1147 1148 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1170 1171 1172 1173 1174 1175 1176 1177 1178 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200 1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260 1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290 1291 1292 1293 1294 1295 1296 1297 1298 1299 1300 1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320 1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350 1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380 1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400 1401 1402 1403 1404 1405 1406 1407 1408 1409 1410 1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440 1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470 1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 1488 1489 1490 1491 1492 1493 1494 1495 1496 1497 1498 1499 1500 1501 1502 1503 1504 1505 1506 1507 1508 1509 1510 1511 1512 1513 1514 1515 1516 1517 1518 1519 1520 1521 1522 1523 1524 1525 1526 1527 1528 1529 1530 1531 1532 1533 1534 1535 1536 1537 1538 1539 1540 1541 1542 1543 1544 1545 1546 1547 1548 1549 1550 1551 1552 1553 1554 1555 1556 1557 1558 1559 1560 1561 1562 1563 1564 1565 1566 1567 1568 1569 1570 1571 1572 1573 1574 1575 1576 1577 1578 1579 1580 1581 1582 1583 1584 1585 1586 1587 1588 1589 1590 1591 1592 1593 1594 1595 1596 1597 1598 1599 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619 1620 1621 1622 1623 1624 1625 1626 1627 1628 1629 1630 1631 1632 1633 1634 1635 1636 1637 1638 1639 1640 1641 1642 1643 1644 1645 1646 1647 1648 1649 1650 1651 1652 1653 1654 1655 1656 1657 1658 1659 1660 1661 1662 1663 1664 1665 1666 1667 1668 1669 1670 1671 1672 1673 1674 1675 1676 1677 1678 1679 1680 1681 1682 1683 1684 1685 1686 1687 1688 1689 1690 1691 1692 1693 1694 1695 1696 1697 1698 1699 1700 1701 1702 1703 1704 1705 1706 1707 1708 1709 1710 1711 1712 1713 1714 1715 1716 1717 1718 1719 1720 1721 1722 1723 1724 1725 1726 1727 1728 1729 1730 1731 1732 1733 1734 1735 1736 1737 1738 1739 1740 1741 1742 1743 1744 1745 1746 1747 1748 1749 1750 1751 1752 1753 1754 1755 1756 1757 1758 1759 1760 1761 1762 1763 1764 1765 1766 1767 1768 1769 1770 1771 1772 1773 1774 1775 1776 1777 1778 1779 1780 1781 1782 1783 1784 1785 1786 1787 1788 1789 1790 1791 1792 1793 1794 1795 1796 1797 1798 1799 1800 1801 1802 1803 1804 1805 1806 1807 1808 1809 1810 1811 1812 1813 1814 1815 1816 1817 1818 1819 1820 1821 1822 1823 1824 1825 1826 1827 1828 1829 1830 1831 1832 1833 1834 1835 1836 1837 1838 1839 1840 1841 1842 1843 1844 1845 1846 1847 1848 1849 1850 1851 1852 1853 1854 1855 1856 1857 1858 1859 1860 1861 1862 1863 1864 1865 1866 1867 1868 1869 1870 1871 1872 1873 1874 1875 1876 1877 1878 1879 1880 1881 1882 1883 1884 1885 1886 1887 1888 1889 1890 1891 1892 1893 1894 1895 1896 1897 1898 1899 1900 1901 1902 1903 1904 1905 1906 1907 1908 1909 1910 1911 1912 1913 1914 1915 1916 1917 1918 1919 1920 1921 1922 1923 1924 1925 1926 1927 1928 1929 1930 1931 1932 1933 1934 1935 1936 1937 1938 1939 1940 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950 1951 1952 1953 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## INTERNATIONAL SEARCH REPORT

Int. Application No

PCT/US 99/12253

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N9/64 C12N15/57 C07K14/47 C07K16/40 C12Q1/68  
A61K39/395

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	PAOLONI-GIACOBINO A. ET AL., : "Cloning of the TMPRSS2 gene, which encodes a novel serine protease with transmembrane, LDLRA, and SRCR domains and maps to 21q22.3." GENOMICS, vol. 44(3), 1997, page 309-320 XP000856785 cited in the application the whole document	1-5, 16, 17
X	DATABASE EMBL NUCLEOTIDE AND PROTEIN SEQUENCES, 11 March 1997 (1997-03-11), XP002132696 HINXTON, GB AC = AA244195. Homo sapiens cDNA clone IMAGE:1007327 similar to SW:KAL_RAT P14272 PLASMA KALLIKREIN PRECURSOR. EST. abstract	1, 3, 16, 17
	— — — — — -/-	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

10 March 2000

Date of mailing of the international search report

23/03/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Mateo Rosell, A.M.

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/12253

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL NUCLEOTIDE AND PROTEIN SEQUENCES, 20 May 1997 (1997-05-20), XP002132697</p> <p>HINXTON, GB</p> <p>AC = AA423272. Mus musculus cDNA clone 820366 5' similar to SW:KAL_HUMAN P03952 PLASMA KALLIKREIN PRECURSOR ;EST. abstract</p>	1, 3, 16, 17
A	<p>WO 94 16083 A (GENETICS INST)</p> <p>21 July 1994 (1994-07-21)</p> <p>page 4, line 7-11</p> <p>page 28-30</p>	1, 3
A	<p>FAIR W R ET AL., : "Prostate-specific membrane antigen."</p> <p>PROSTATE,</p> <p>vol. 32 (2), 1997, page 140-148</p> <p>XP000870112</p> <p>the whole document</p>	5-15
A	<p>WO 93 01309 A (SCRIPPS RESEARCH INST)</p> <p>21 January 1993 (1993-01-21)</p> <p>abstract</p> <p>page 64, line 20 -page 66, line 21</p>	6-15
A	<p>YOUNG C Y F ET AL., : "Expression and androgenic regulation of human prostate-specific kallikreins"</p> <p>JOURNAL OF ANDROLOGY,</p> <p>vol. 16, no. 2, April 1995 (1995-04), pages 97-99, XP000856769</p> <p>cited in the application</p> <p>the whole document</p>	1
A	<p>TANIMOTO H T AL., : "Hepsin, a cell surface serine protease identified in hepatoma cells, is overexpressed in ovarian cancer"</p> <p>CANCER RESEARCH,</p> <p>vol. 57 (14), 1997, page 2884-2887</p> <p>XP000867297</p> <p>cited in the application</p> <p>the whole document</p>	1
A	<p>YAMAMOTO M ET AL., : "Raised prostate-specific antigen in adenocarcinoma of the colon"</p> <p>INTERNATIONAL UROLOGY AND NEPHROLOGY ,</p> <p>vol. 29 (2), 1997, page 221-225</p> <p>XP000889898</p> <p>the whole document</p>	6, 7, 37
	-/-	

## INTERNATIONAL SEARCH REPORT

Inte Application No

PCT/US 99/12253

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	<p>DATABASE EMBL NUCLEOTIDE AND PROTEIN SEQUENCES, 16 November 1998 (1998-11-16), XP002132700 HINXTON, GB AC = AI261741. Homo sapiens cDNA clone IMAGE:2028487 3' DE similar to TR:015393 015393 SERINE PROTEASE. ;, mRNA sequence. abstract</p> <hr/>	1, 3

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/12253

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9416083 A	21-07-1994	AU 675506 B	06-02-1997
		AU 6089894 A	15-08-1994
		CA 2153254 A	21-07-1994
		EP 0679189 A	02-11-1995
		JP 8502176 T	12-03-1996
		US 5665566 A	09-09-1997
WO 9301309 A	21-01-1993	US 5288612 A	22-02-1994
		AT 160876 T	15-12-1997
		AU 652474 B	25-08-1994
		AU 2317492 A	11-02-1993
		DE 69223355 D	15-01-1998
		DE 69223355 T	28-05-1998
		DK 528525 T	23-02-1998
		EP 0528525 A	24-02-1993
		EP 0595865 A	11-05-1994
		ES 2109982 T	01-02-1998
		GR 3025752 T	31-03-1998